# Influence of Date of Sowing on Growth and Yield of Different Varieties of Radish (*Raphanus Sativus Linn.*) Under Allahabad Agro Climatic Conditions

## Evakordor K. Jyrwa<sup>1</sup>, Dr. B. Mehera<sup>2</sup>

<sup>1</sup>Department of Environmental Sciences

<sup>2</sup>Sam Higginbottom University of Agriculture Technology and Sciences, Allahabad, U.P

Abstract: An experiment was conducted at the Research farm, School of Forestry and Environment, SHIATS, Allahabad during Rabi season 2014-2015 to study the effect of sowing dates on three different varieties of radish. The experiment was laid out in Randomized Block Design (3x3 factorial) and replicated thrice. Seeds of the cultivars viz., hill queen; Mino early and S.S11 were sown on three different dates i.e. Oct 7, Oct 17 and Oct 27. Maximum plant height, leaf length, number of leaves, root length and root yield was obtained by 3rd sowing i.e. Oct 27. While root diameter and root weight were obtained by 1st sowing i.e. Oct 7. It was observed that Mino early recorded the highest root yield about 13.10 t/ha. The combined effect of sowing dates and varieties showed that sowing on Oct 27, all varieties performed well in respect of yield and yield components. Mino early was found to be the best in respect of quality judging.

Keywords: Radish spp., root yield, Mino early, sowing time

## 1. Introduction

Radish (Raphanus sativus Linn.) is an annual or biennial herb depending on types. It is a fast-maturing, easy-to-grow root vegetable crop grown in both tropical and temperate regions. Radish is a good source of vitamin C (ascorbic acid), containing 15-40mg/100g of edible portion and contains a variety of minerals. It is very difficult to raise the good quality radish roots because of excessive forking, splitting, cracking and pithiness, which is highly influenced by varieties and time of sowing. Hence, selection of optimum sowing time, good quality varieties and spacing's are key factors for successful radish production

Arora and Pandey (1969) reported that radish sown between the end of September and mid-November yielded more than that sown later. The best sowing time in South India is from April–June in the hills and October-December in the plains (**Purewal**, 1957). The average yield of Indian cultivars varies from 150-200 quintals/ha.

Proper sowing time depend on the varieties and prevailing environment. Therefore the selection of right type of varieties for sowing at optimum time is the key factor for successful radish production. Growers tend to manipulate sowing time in order to obtain better growth and higher quality yield. The time of sowing is also adjusted so as to synchronize the time of harvest with market demand.

Under this circumstance, the present study was undertaken with the aim of investigating the effects of sowing date on growth and yield of three varieties of radish. It is anticipated that the information gathered from the results of the present experiment would help the growers to increase the production of radish with favorable weather conditions and proper sowing time.

### 2. Material and methods

The field experiment was carried out at the Research farm, School of Forestry and Environment, SHIATS-Deemed University, Allahabad during the period from October 2014-December 2014. The soil of the experiment was sandy loam in texture. The land was well drained with good irrigation facilities. The experiment was designed to study the influence of sowing date (D1, October 7th; D2, October 17th and D<sub>3</sub>, October 27<sup>th</sup>) on growth and yield of three varieties (V<sub>1</sub>, hill queen; V<sub>2</sub>, Mino early; V<sub>3</sub>, S.S11) of radish. Two factorial experiment was laid out following a Randomized block design (R.B.D) with three replications. The whole experimental area was 144m2 each block was divided into nine plots where nine treatments were allotted at random. The size of the plot was 2m x 2m, with the spacing of 30cm x 10cm. Well decomposed FYM and a basal dose of fertilizer were applied during final preparation. Irrigation and drainage channel were prepared around the plot before sowing the seeds. Seeds were sown on different sowing times with 10 days interval, in about 1.5cm depths, and in lines continuously and covered by loose soil. Seedling emergence was completed within 7 days after sowing. Seedlings were thinned out and it was done after 15 days of sowing. Weeding was done as or when required to keep the plot free from weed and to pulverize soil. General irrigation was done by twice in a week. The crop was harvested periodically and harvesting was done at 45 days after each sowing date. To evaluate the effect of sowing time on three selected varieties, following observations were made to get information related to plant growth as well as yield. Data were collected from experimental plots on different growth and yield components and yield were statistically analyzed.

### 3. Results and Discussion

The observation recorded on growth and yield of radish vegetable crop is influenced by variety and dates of sowing and observed that the 3<sup>rd</sup> sowing i.e. 27<sup>th</sup> October proved to

# Volume 7 Issue 6, June 2018 www.ijsr.net

Licensed Under Creative Commons Attribution CC BY

### International Journal of Science and Research (IJSR) ISSN (Online): 2319-7064 Index Copernicus Value (2016): 79.57 | Impact Factor (2017): 7.296

be most superior with respect to early germination and among the three dates of sowing maximum time taken to germinate was observed in the first sowing i.e. 7th October whereas among the varieties the 2<sup>nd</sup> variety i.e. mino early observed minimum time taken to germination and found maximum in the 3<sup>rd</sup> variety i.e. the delayed in germination might be due to the increased in minimum temperature in the first sowing since radish is a cool season crop that required a minimum temperature of 18°C during germination. A number of days taken to maturity were highest in the 1st sowing, the increased in days to maturity might be due to longer duration of crop period and lowest in third sowing due to forced maturity caused by an increase in temperature. Seeds sown earlier took more time for the maturity of roots. Among the yield attributes, root length and root yield was observed maximum in the 3<sup>rd</sup> sowing and minimum in the 1<sup>st</sup> sowing whereas among the varieties maximum root length was observed in the 1<sup>st</sup> variety and minimum in 3<sup>rd</sup> variety and maximum root yield was observed in the 2<sup>nd</sup> variety i.e. mino early and minimum root yield was found in the 1<sup>st</sup> variety i.e. hill queen. (**Alam et al, 2010**) he suggested that sowing of radish variety from the first fortnight of October to November gave the longest root length. The higher yield in above treatments is due to better plant survival owing to the favorable environmental conditions for growth and development of roots and the closer spacing accommodates a number of plants per unit area. Similar results were reported by **Rehman and Nawab Ali (2000).** 

And the effect of interaction between dates of sowing and varieties differed significantly for all the yield and growth parameters with a p-value ( $p \ge 0.05$ ) as shown in Tables 1&2. It shows that maximum root yield was obtained by mino early sown during the late October i.e.  $27^{th}$ 

Table 1: Effect of sowing dates and varieties on days to germination and days to maturity

	Days to germination				Day						
Date of sowing	Variety										
	$V_1$	$V_2$	V <sub>3</sub>	Mean	<b>V</b> <sub>1</sub>	$V_2$	V <sub>3</sub>	Mean			
$D_1$	7.133	6.667	7.133	6.977	54.333	55.333	57.000	55.556			
$D_2$	6.100	6.300	6.200	6.200	53.000	48.637	50.067	50.578			
D <sub>3</sub>	5.967	5.700	5.933	5.866	45.733	46.111	40.000	43.956			
Mean	6.400	6.222	6.422		51.022	50.044	49.022				
Interaction effect											
Source	D	V	DV		D	V	DV				
S.E(d)	0.181	0.181	0.314		1.955	1.955	3.387				
C.D	0.384	0.384	0.666		4.146	4.146	7.181				

 Table 2: Effect of sowing dates and varieties on root length (cm) and root yield (t/ha)

Date of sowing	Root length (cm)				Root yield (t/ha)					
	Variety									
	V <sub>1</sub>	$V_2$	$V_3$	Mean	V <sub>1</sub>	$V_2$	$V_3$	Mean		
$\mathbf{D}_1$	22.333	20.333	18.667	20.444	9.797	12.00	11.183	10.993		
$\mathbf{D}_2$	22.900	20.567	19.567	21.011	10.810	11.733	10.533	11.026		
$D_3$	24.133	22.500	20.067	22.233	10.133	13.100	10.680	11.301		
Mean	23.122	21.133	19.433		10.240	12.281	10.799			
Interaction effect										
Source	D	V	DV		D	V	DV			
S.E(m)	1.376	1.376	2.383		0.643	0.643	1.114			
C.D	1.180	1.180	2.044		1.364	1.364	2.363			

## 4. Conclusion

The experimental results were observed maximum root yield (13.10t/ha) in treatment combination  $D_3V_2$  (October 27 and Mino early) due to maximum plant survival and maximum plant population per unit area. Therefore based on the experiment performed we can be concluded that sowing of radish between late of October and the first week of November gave maximum root yield.

Hence, the radish variety Mino early and sowing date October 27th was most suitable for Allahabad Agro-climatic conditions.

## References

[1] Ahmed, M. J. and Siddique (2004). Effect of sowing dates on growth and yield of broccoli (*Brassica oleracea L.*). Asian J. Plant Sci., 3(2): 167-169.

- [2] Alam, M., Farooque, A.M., Nuruzzaman, M. and Jamal Uddin (2010). Effect of sowing time on growth and yield of three radishes (*Raphanus sativus L.*) varieties. *Bangladesh Res Pub J* 3(3): 998-1006.
- [3] Arora, P.N and Pandey, S.L (1971). Effect of time of sowing radish on the yield economic returns. *Hort. Abstr.* 41(1): 137
- [4] Aziz-Ur-Rehman and Ali (2000). Effect of plant spacing and sowing time on yield in turnip (*Brassica* campestris cv. Purple Top). Sarhad J. Agri., 16(6): 575-579.
- [5] **Purewal, S.S. (1957).** Vegetable Cultivation in North India, Farm Bull., *Indian Council of Agricultural Research*, New Delhi, No. **36**, p. 24.
- [6] Pervez, M. A., Ayub, C.M., Saleem, B.A., Virk, N.A. and Mahmood, N. (2004). Effect of nitrogen levels and spacing on growth and yield of radish (*Raphanus* sativus L.). Int. J. Agri. Biol., 6(3): 504-506.

<u>www.ijsr.net</u>

#### Licensed Under Creative Commons Attribution CC BY