

Congenital Gastrointestinal Anomalies and their Associations to Genetic Disorders

Aurel Vula¹, Alba Dokaj²

^{1,2} Department of Pediatrics, University Hospital Centre "Mother Teresa", Tirana, Albania

Abstract: *Congenital anomalies of the digestive tract are an important part of congenital anomalies as they constitute one of the main causes of mortality in children. The aim of this study was to evaluate the most common digestive anomaly associated with genetic disorders. We also sought to ascertain the most frequent genetic disorder in our patients and its connection to mortality, the average age of diagnosis, prenatal diagnosis, gestational age and associations with other organ anomalies. The study was conducted in two phases and includes a total of 273 children diagnosed with Congenital anomalies of the digestive tract, presented at the Pediatric Intensive Care Unit in UHC "Mother Teresa". The first phase, retrospective, includes 137 patients during the period January 2006 – December 2010. The second phase of the study, prospective, includes 136 patients from January 2011 - March 2015. The most frequent digestive anomaly in patients with genetic disorders was Anal atresia(39.40%), followed by Omphalocele and Intestinal atresia (15.10%). Down syndrome was the most frequent genetic disorder, 9 cases (3.30%). It is important to identify cases associated to genetic disorders because they condition patient's management and prognosis. Anomalies that are often associated with other defects have shown early clinical symptoms and require more time and effort for correction.*

Keywords: congenital gastrointestinal anomalies, newborn babies, genetic disorders, age of diagnosis, associated organ anomalies.

1. Introduction

The gastrointestinal tract may be subject to a variety of congenital abnormalities (i.e. those present at birth) that arise during embryological development. Specific patterns of malformations of the gastrointestinal tract include abnormal lumenisation (stenoses and atresias), duplications, abnormal rotation and fixation, abdominal defects and a variety of others associated with persistence of embryonic structures (e.g. Meckel's diverticulum), or abnormal formation of specific regions of the gastrointestinal tract (e.g. microgastria) or its cellular components (e.g. nerves in Hirschsprung's disease). These disorders primarily result in symptoms of intestinal obstruction, effects on surrounding structures or of associated anomalies.[1]

They frequently manifest with feeding difficulties, distention, and emesis at birth or within 1 or 2 days.[2] The clinical symptoms varies depending on the pathology, the age of diagnosis and associated anomalies. They may be from mild to severe with, recurrent abdominal pain, intestinal obstruction, dehydration, malnutrition, malabsorption/diarrhea, peritonitis/septic shock, solid food intolerance, common bile duct obstruction, abdominal distention, and failure to thrive.[3] Some congenital GI malformations, such as malrotation, have a very good outcome, whereas others, such as congenital diaphragmatic hernia, have a poor outcome, with a relatively high mortality rate of 10 to 30%. [2]

Early clinical recognition of these disorders is essential to minimise complications and allow the early institution of appropriate therapies.

Although surgery is the commonest intervention early treatment must include adequate resuscitation and stabilisation of the child prior to definitive surgery. Many of these abnormalities are associated with other congenital

anomalies or genetic syndromes and disease making the clinical symptoms and the management more complex and difficult. [1]

A genetic disorder is a disease caused in whole or in part by a change in the DNA sequence away from the normal sequence. Genetic disorders can be caused by a mutation in one gene (**monogenic disorder**), by mutations in multiple genes (**multifactorial inheritance disorder**), by a combination of gene mutations and environmental factors, or by damage to chromosomes, changes in the number or structure of entire chromosomes, the structures that carry genes (**chromosome disorders**).[4]

The incidence of disease syndromes and genetic defects in patients with malformations is much higher than in the general population.[5] Since genetic syndrome is seen to have an important impact in patient's life, in the U.S. and many other countries there are created healthcare programs for newborn genetic screening that identify treatable genetic disorders in newborn infants. Early intervention to treat these disorders can eliminate or reduce symptoms that might otherwise cause a lifetime of disability.[6]

Through our study we sought to evaluate the connection between genetic disorders and congenital gastrointestinal anomalies in newborns who were hospitalized in our Intensive Unit Care.

2. Material and methods

The study was conducted in two phases and includes a total of 273 children diagnosed with Congenital anomalies of the digestive tract, presented at the Pediatric Intensive Care Unit in UHC "Mother Teresa". The first phase of the study is case-control type or otherwise known as retrospective study. It includes 137 patients, who were admitted to the Pediatric Intensive Care Unit during the period January 2006 –

December 2010 with the main diagnose "Congenital gastrointestinal anomalies".

The second phase of the study is prospective type and includes cases presented from January 1, 2011 to March 1, 2015, where patients have been followed for at least 1 month. The second phase of the study, prospective, includes 136 patients.

Statistical analysis

Continuous data were presented in average value and in standard deviation. Discrete data were presented in absolute value and in percentage. The correlation between the two variables was analyzed by the Kendal's tau correlation coefficient.

Presentation of the data was done through simple and composite tables and graphs of different types.

The statistical analysis was carried out through the statistical package SPSS 19.0 (Statistical Package for Social Sciences Inc., Chicago II USA) and Microsoft Excel.

Significant values of $p \leq 0.01$ were considered.

3. Results and discussion

In our study are included 273 patients with Congenital digestive anomaly as above in table 1:

Table 1: Congenital digestive anomaly frequency

Congenital digestive anomaly	No.	(%)
Anal atresia	59	21.60%
Omphalocele	38	13.90%
Intestinal atresia	50	18.30%
Meconium ileus	3	1.10%
Hirschsprung's disease	8	3.00%
Duodenal atresia	20	7.30%
Esophageal atresia	48	17.60%
Biliary atresia	6	2.20%
Diaphragmatic hernia	26	9.50%
Intestinal malrotation	3	1.10%
Common mesentery	6	2.20%
Congenital megacolon	2	0.70%
Hypertrophic pyloric stenosis	4	1.50%
Total	273	100.00%

The most frequent anomaly in our study is anal atresia (21.60%), followed by intestinal atresia (18.3%) and esophageal atresia (17.60%).

In a total of 273 cases diagnosed with Congenital gastrointestinal anomaly, 33 of them (12.10%) had also a genetic syndrome or disease.(table 2)

Table 2: Frequency of genetic disorders in our patients

Genetic disorder	No.	(%)
Present	33	12.10%
Not present	240	87.90%
Total	273	100%

From 33 cases presents with genetic disorder the most common digestive anomaly was found Anal atresia (39.4%). In our study biliary atresia, diaphragmatic hernia, intestinal malrotation, common mesentery, congenital megacolon and hypertrophic pyloric stenosis have not been associated with genetic disorders. (table 3)

Table 3: Frequency of digestive anomalies associated with genetic disorders

Congenital digestive anomaly	No.	(%)
Anal atresia	13	39.40%
Omphalocele	5	15.10%
Intestinal atresia	5	15.10%
Meconium ileus	3	9.10%
Hirschsprung's disease	3	9.10%
Duodenal atresia	2	6.10%
Esophageal atresia	2	6.10%
Biliary atresia	0	0%
Diaphragmatic hernia	0	0%
Intestinal malrotation	0	0%
Common mesentery	0	0%
Congenital megacolon	0	0%
Hypertrophic pyloric stenosis	0	0%
Total	33	100.00%

The most frequent genetic disorder seen in our study was Down syndrome, with 9 cases (3.30%), followed by Cystic fibrosis, 5 cases (1.80%). (table 4)

Table 4: Genetic disorders by their frequency

Genetic disorder	No.	(%)
S.Down	9	3.30%
Cystic fibrosis	5	1.80%
VACTERL	1	0.37%
OEIS Complex	1	0.37%
Trisomy 18	1	0.37%
Noonan	1	0.37%
Without genetic disorder	240	87.90%
Others	15	5.50%
Total	273	100%

In our study was found a significative relation ($p < 0.01$) between genetic disorders and other associated anomalies in patients with congenital digestive anomalies. Most frequent associated anomalies were found heart (20.50%), pulmonary (7.70%), abdominal (5.50%) and renal diseases (4.0%). 23 (69.70%) of 33 cases with Congenital digestive anomaly and genetic disorder, also had another associated anomaly. (table 5). In another study made in Hospital Infanta Cristina, SES, Spain was considered that it is important to identify any associated anomalies, especially heart disease, craniofacial anomalies and other gastrointestinal malformations, because they condition the patient's management. The knowledge of these patients is essential for correct treatment. [5]

Tabela 5: Associated anomalies

Genetic disorder	Associated anomalies	No other anomaly	Total
Present	23	10	33
	69.70%	30.30%	100%
Not present	97	143	240
	40.40%	59.60%	100%

Also data from the literature showed that children with congenital heart disease have a much higher incidence of intestinal malformations than those with normal heart and that they frequently present with multiple malformations (chromosome aberrations or multiple organ lesions). This multiple malformation complex is particularly common in anorectal malformations where the incidence of congenital heart diseases is 9 to 14%, with predominance of VSD and tetralogy of Fallot. [7]

Another significant connection ($p < 0.01$) was found between the presence of genetic disorders and the diagnostic age of patients with congenital digestive anomalies. The more associated anomalies, the more symptoms are presented, and this is reflected in the age of diagnosis. In cases with genetic disorder the average age of diagnosis was about two times smaller than in cases with isolated congenital digestive anomaly.(table 6)

Table 6: Connection with average age of diagnosis

Genetic disorder	Average diagnostic age
Present	2.24 days
Not present	4.3 days

Between mortality rate, prenatal diagnosis, gestational age, birth weight and genetic disorders was not found a significant correlation ($p > 0.01$).

4. Conclusion

Identification of patients with genetic syndrome or disease plays a fundamental role in management and prognosis of patients diagnosed with Congenital digestive anomalies. Coexistence with multiple anomalies makes the clinic presentation more severe and more complex to achieve a better management.

References

- [1] Fotis, Lampros, Burns, Alan J, and Thapar, Nikhil(Aug 2012) Gastrointestinal Tract: Congenital Abnormalities. In: eLS. John Wiley & Sons Ltd, Chichester. <http://www.els.net> [doi: 10.1002/9780470015902.a0002138.pub2]
- [2] Merck Sharp & Dohme Corp., a subsidiary of Merck & Co., Inc., Kenilworth, NJ, USA
- [3] Biliary vomiting in the newborn: Rapid diagnosis of intestinal obstruction. Ken Kimura, M.D., and Vera Loening-Baucke, M.D., University of Iowa College of Medicine, Iowa City, Iowa. *Am Fam Physician*. 2000 May 1;61(9):2791-2798
- [4] www.genome.gov/19016930/faq-about-genetic-disorders/
- [5] Digestive malformation and their associations to syndrome condition and genetic defects, Enríquez Zarabozo E, Blesa Sánchez E, Ayuso Velasco R, Galán Gómez E
- [6] learn.genetics.utah.edu/content/disorders/screening/
- [7] Intestinal malformation and congenital heart diseases. Voisin M, Galifer RB, Kadiri T, Grolleau R, Dumas

R, Jean R. Arch Mal Coeur Vaiss. 1987 Apr;80(4):524-8.

Author Profile



Aurel Vula, Pediatrician Division of General Pediatrics, Pediatric Rheumatology and Cardiology, Department of Pediatrics, University Hospital Centre "Mother Teresa", Tirana, Albania



Alba Dokaj, Pediatric Resident, Department of Pediatrics, University Hospital Centre "Mother Teresa", Tirana, Albania