Antifungal Potentiality of Leaf Extracts of Saraca indica Against Fusarium Species, An Approach towards Eco-friendly Management

Pushpa Kumari¹, Reena Mohanka², Priyanka³

^{1, 2, 3}Plant Pathology and Microbiology Laboratory, Department of Botany, Patna University, Patna-800005, India

Abstract: Fusarium species are the most frequently soil borne fungal pathogen on vegetable crops, causing significant yield losses and made economical problems for growers. Fusarium control through chemical fungicides causes serious environmental problems and are toxic to non-target organisms as well. Plant based pesticides appear to be one of the better alternatives as they are known to have minimal environmental impact and danger to consumer. In our study, in-vitro antifungal potentiality of leaf extract (aqueous and ethanolic) of Saraca indica was tested against Fusarium solani and Fusarium moniliforme through poisoned food technique and was compared with leaf extracts of Azadirachta indica for five different concentrations(10%, 20%, 30%, 40% and 50%). Ethanolic extract of Saraca indica was found to be more efficient showing excellent inhibitory effect, 77.22% for Fusarium solani and 80% for Fusarium moniliforme at 50% concentration, in comparison to that of Azadirachta indica,70.83% and 72.77% respectively. The results indicate that some common plants could be exploited in developing a potent plant based fungicides which can be used in organic farming for the eco-friendly management of fusarium species.

Keywords: Leaf extracts, Fusarium solani, Fusarium moniliforme, Azadirachta indica, Poisoned food technique

1. Introduction

Among the greatest hazards in crop-production, diseases and pests are the main problems. Fusarium species are the best known soil borne plant pathogens in terms of economical damage in agricultural productions all over the world[1] -[3] .Dry rot on potato, wilting and decline on bean or pea, crown rot and head blight on wheat, bakanae disease on rice caused by Fusarium species result to yield losses in most crop fields[4].Fusarium solani (Mart.) Sacc., a soil inhabiting pathogen, attacks a large number of host plants, including oilseeds pulses, vegetables and ornamentals [5] -[8].Fusarium dry rot of seed tuber can reduce crop establishment by killing developing potato, where crop losses can be up to 25%, while more than 60% of tubers can be infected in storage [9]. Many methods such as chemical, cultural and biological techniques have been developed for the control of plant disease by soil borne pathogens [10]. The efforts were made in other regions to manage the Fusarium species through phytoextracts [11] - [15] in various crops.

Saraca indica, commonly known as Ashoka and one of the most common tree of India, is an evergreen tree with numerous spreading and drooping glabrous branches and has been extensively used in Ayurveda. This plant has shown many pharmacological properties such as antidiabetic [16], antiulcer [17], anthelmintic [18] and antimicrobial [19] – [20] activities.

The present study is designed to evaluate the antifungal properties of Ashoka tree against destructive soil borne plant pathogenic fungus *Fusarium* and also the percentage inhibition is compared with *Azadirachta indica*(Neem- a well known antibacterial and antifungal agent).

2. Literature Survey

Azadirachta indica, commonly known as a Neem, found throughout India is the most extensively used tree in the field of medicine. It is also used world-wide to control different plants diseases and pests. Study shows that growth of soil borne pathogens (*Fusarium oxysporumf.* sp. cicero, *Rhizoctonia solani, Sclerotium rolfsii* and *Sclerotinia sclerotiorum*) in liquid medium was inhibited by extracts of leaf, trunk, bark, fruit pulp and oil of neem[21]. Aqueous leaf extracts of neem exhibited considerable control of *Fusarium oxysporum* in banana [22].

Saraca indica is also a massive tree found throughout India and used in Ayurveda. The bark of this plant is used as astringent, demulcent, emollient and stomachic. The leafis used to treat stomachalgia and flowers are used for treating of syphilis, inflammation, dysentery, haemorrhoids and scabies in children [23]. It is also screened to control microbes (fungi and bacteria) causing different plant diseases leading to a great crop loss. This paper screens the antifungal properties of leaf extracts *Saraca indica* against some soil borne pathogens escpecially *Fusarium* species to be used as fungicide next to *Azadirachta indica*.

3. Material and Methods

Plant materials

Fresh leaves of two plants, *Azadirachta indica* and *Saraca indica* were collected from local area of Patna. The plants were authenticated by taxonomist from Botany Department, P.U., Patna.

Extract preparation

Collected fresh leaves of the aforesaid plants were washed separately with tap water followed by sterile distilledwater and dried in shade. The dried materials were finally grinded

DOI: 10.21275/ART20183883

to powder, sealed in polythene bags and stored away from light and moisture until used for extraction[24] - [26].

Aqueous extract- 50gm of each material was soaked in 200ml of distilled water for 30 minutes and then boiled to half volume. After cooling, filtered with muslin cloth followed by Whatman filter paper no.-1 and kept in dark glass bottles at 5 $^{\circ}$ C in a refrigerator.

Ethanolic extract-50gm of each material was homogenized in 200ml mixture ofethanol anddistilled water (50:50, v:v) and left in dark bottlesfor72hrs on shaker.Then extracts were filtered with muslin cloth, followed by Whatman filter paper no.-1in other dark bottles and exposed to 60 °C in water bath for 30 minutes for ethanol evaporation. Extracts were stored in dark bottles at 5°C.

Isolation of pathogen

Fusarium species were isolated through serial dilution culture technique of soil collected from vegetable crop fields. PDA(Potato Dextrose Agar) was used as growth medium. Further a pure culture of each colony type, growing in petri plates, was obtained and maintained by subculturing. The temperature was maintained at 25 ± 2 °C. Cultures were identified on the basis of macro and microscopic characteristics, reverse surface coloration of colonies, conidial morphology and slide culture technique[27] – [31]. The technique of slide culture was also used to identify the species of *Fusarium*[32] which allow the direct microscopic observation of morphological structure of taxonomic value.

Identification of *Fusarium* species as *Fusarium solani* and *Fusarium moniliforme* was also confirmed by IARI, New Delhi.



Figure 1: Different fungal-colonies growing in PDA (serial dilution culture of soil)



Figure 2: Isolation of *Fusarium* species from petri plates of soil culture.

Antifungal screening

Different concentration viz-10%, 20%, 30%, 40% and 50% of leaf extracts (both aqueous and ethanolic) of aforesaid plants were screened for their antifungal activities through poisoned food technique [33].Proportionate amount of leaf extracts of two botanicals were added to separate flasks containing media to prepare media having 10% -50% concentration of leaf extracts. To avoid bacterial contamination an antibiotic (Chloramphenicol, 0.10 mg/l) was supplemented to the media.20 ml of media was poured into petri plates, allowed to solidify and inoculated individually with 5mm diameter discs of the tested F. solani and F. moniliforme. Plates with media, not supplemented with leaf extracts and inoculated with Fusarium species, served as negative control. After seven days of incubation at $25 \pm 2^{\circ}$ C, orthogonal measurements of colonies were taken using the control plates as a reference. The percentage inhibition of growth was calculated according to following formula-

% inhibition = (dc-dt/dc) 100

Where, dc=Average increase in mycelial growth in control. dt= Average increase in mycelia growth in treatment [34].

The two botanicals were also screened for antifungal activities in combination of 10% and 20% each and the two combination with a fungicide in a very small percentage (Bebestin, 0.5% of mother solution-5gm/l).



Figure 3: Comparative study of the inhibitory effect of leaf extracts of *A. indica* (*A*) and *S. indica* (*B*) on *Fusarium*



Figure 4aFigure 4bFigure 4: Growth of Fusarium in 50% alcoholic extract of
A. indica (4a) and S. indica (4b)

4. Results and Discussion

Results presented in Table 1, 2, 3, 4 and 5 show the inhibitory effect of aqueous (Table-1 and3) and ethanolic (Table-2, 4 and 5) leaf extracts of the two botanicals against Fusarium species. Effects of the leaf extracts of the two botanicals were shown in Table 1 and 2 on Fusarium solani and in Table 3 and 4 on Fusarium moniliforme. For both the plant extracts radial growth of tested fungi decreased significantly with increased concentration of added extracts. The interesting observation was that ethanolic extracts were found more effective then aqueous one and also Saraca indica was found more effective in both the conditions. Also the botanical control of Fusarium moniliforme was more than Fusarium solani, as 50% ethanolic leaf extract concentration of Saraca indicainhibited 80% of the mycelial growth of Fusarium moniliforme (upto 77% mycelial growth inhibition of Fusarium solani). The result could be used as an indicator to exploit Saraca indica, a common medicinally important evergreen tree, as a fungicide.

The two botanicals leaf extracts were also tested for their inhibitory effects in a combination (Table-5). There were four formulations. In first, PDA medium had 10% leaf extracts of each plant and in the second one it had 20% each. The two botanicals showed a superior inhibitory effects on radial fungal growth in their mixed combination. In third and fourth formulations, 0.5% of Bebestin (a fungicide) was supplemented to aforesaid first and second formulations and these showed 100% control of *Fusarium* species in 20% leaf extract preparation with Bebestin (fourth formulation). Even in 10% each botanical leaf extracts + 0.5% Bebestin combination, there was a 100% inhibition for *Fusarium moniliforme* and 77.78% for *Fusarium solani* which was also a remarkable result.

 Table 1: Inhibitory effect of aqueous leaf extracts of

 Azadirachta indica and Saraca indica on mycelial growth of

 Eusarim solani

Tusa III solali				
	Azadirachta indica		Saraca indica	
Different	Mycelial	%	Mycelial	%
concetration of	growth in	Inhibition	growth in	Inhibition
aqueous leaf extracts	mm		mm	
10%	82.5	8.33	81	10
20%	77.25	14.16	77	14.44
30%	66.5	26.11	63	30
40%	56	37.77	35.5	40.55
50%	44	51.11	38	57.77



Figure 5: Graphical representation of Table-1

 Table 2: Inhibitory effect of ethanolic leaf extracts of

 Azadirachta indica and Saraca indica on mycelial growth of

 Fusarim solani

i usainii solan					
	Azadirachta indca		Saraca indica		
Different	Mycelial	%	Mycelial	%	
concentration of	growth	Inhibition	growth in	Inhibition	
ethanolic leaf extracts	in mm		mm		
10%	77.5	13.88	74	17.77	
20%	68.0	24.44	64	28.88	
30%	58.5	35.00	53.25	40.83	
40%	37	58.88	34.5	61.66	
50%	26.25	70.83	20.5	77.22	



Figure 6: Graphical representation of Table-2

 Table 3: Inhibitory effect of aqueous leaf extracts of

 Azadirachta indica and Saraca indica on mycelial growth of

Fusarium moniliforme				
	Azadirachta indica		Saraca indica	
Different	Mycelial	%	Mycelial	%
concentration of	growth in	Inhibition	growth in	Inhibition
aqueous leaf extracts	mm		mm	
10%	81	10	78	13.33
20%	73.5	18.33	72	20
30%	63.5	29.72	62	31.11
40%	50	44.44	48.5	46.11
50%	41	54.44	36	60

DOI: 10.21275/ART20183883



Figure 7: Graphical representation of Table-3

Table 4: Inhibitory effect of ethanolic leaf extracts of Azadirachta indica and Saraca indica on mycelial growth of Eusarium moniliforme.

i usurum mommorme.					
	Azadirachta indica		Saraca indica		
Different concentration	Mycelial	%	Mycelial	%	
of ethanolic leaf	growth	Inhibition	growth	Inhibition	
extracts	in mm		in mm		
10%	70	22.2	63	30	
20%	60	33.33	51.25	43.06	
30%	48.5	46.11	44.25	50.83	
40%	36	60	30.5	66.11	
50%	24.5	72.77	18	80	



Figure 8: Graphical representation of Table-4

 Table 5: Inhibitory effect of ethanolic leaf extracts of

 Azadirachta indica and Saraca indica on mycelial growth of

 Fusarium solani and Fusarium moniliforme

r usarium soram and r usarium monimorme					
	Fusarium solani		Fusarium moniliforme		
Concentrati-on of leaf	Mycelial	%	Mycelial	%	
extracts of A. indica	growth in	Inhibition	growth in	Inhibition	
and S. indica	mm		mm		
10% of each	63	30	53.25	40.83	
20% of each	36	60	31.5	65	
10% of each +0.5% Bebestin	20	77.78	00	100	
20% of each +0.5% of Bebestin	00	100	00	100	



Figure 9: Graphical representation of Table-5 for leaf extract combination



Figure 10: Graphical representation of Table-5 for leaf extract combination with 0.5% Bebestin

5. Conclusion

This study was focused on screening the antifungal activity in some common botanicals to use as natural, harmless fungicide, aiming at partial or total replacement of chemical synthetic fungicide to control soil borne diseases in vegetable crops. A chemical fungicide, Bebestin, at very low percentage (0.5% of mother solution 5gm/ l) show 100% control in combination with 20% ethanolic extracts of both plants (Table-5).

6. Future Scope

The results obtained in the present study reflect light on successful development of safe and cheap compound as antifungal which provide a potent tool to control not only *Fusarium* species but also for controlling other soil borne pathogens and getting success in the avoidance of environmental pollution and side effect of fungicide.

References

- [1] Saremi H, (2000). Plant Diseases Caused by Fusarium Species. Jihad Daneshgahi, Ferdossy Mashhad University, Iran, p. 160.
- [2] Bentley AR, Cromey M G, Farrokhi-Nejad R, Leslie JF, Summerell BA, Burgess LW, (2006). *Fusarium* crown and root rot pathogens associated with wheat and grass

Volume 7 Issue 7, July 2018 www.ijsr.net

Licensed Under Creative Commons Attribution CC BY

DOI: 10.21275/ART20183883

stem bases on the South Island of New Zealand. Australasian Plant Pathol. 35(5): 495-502.

- [3] Bockus WW, Bowden RL, Hunger RM, Morrill WL, Murray TD, Smiley RW, (2007). Compendium of Wheat Diseases and Insects, 3rd Edition. APS Press, St. Paul, MN.
- [4] Saremi H, JAfary H, Ammarlou A, (2007a). Incidence of crown rot disease of wheat caused *Fusarium pseudograminearum* as a new soil borne fungal species in west Iran. Pak. J. Biol. Sci. 10(20): 3606-3612.
- [5] Mani, A. and Sethi, C. L. 1968. Reduced lamina and a typical symptom induced by *Fuserium solani* on chickpea. *Indian Phytopathology*. 38: 542-543.
- [6] Bazalar, C. J. and Delgadi, M. A. 1981. *Fusarium* species associated with the vascular system of cotton (*Gossypium barbadense* L.) under conditions in Piura, Peru. *Fitopathologia***16**(1): 6-15.
- [7] Kumar, R. U., Safeeulla, K. M. and Shetty, H. S. 1983. Seedborne nature and infectivity of *F. solani* (Mart.) Sacc. The causal agent to wilt and fruit rot of brinjal. *Indian National Science Academic Journal*. 49(2): 134-140.
- [8] Kore, S. S. and Mane, A. V. 1992.Dry root rot of kagzilime seedlings caused by *Fusarium solani*. *Journal* of Maharashtra Agricultural University. 17(2): 276-278.
- [9] Wharton PS, Tumbalam P, Kirk WW, (2006). First Report of Potato Tuber Sprout RPT Caused by *Fusarium sambucinum* in Michigan. Plant Disease 90: 1460-1164.
- [10] Singleton L, Mihail DJ, Rush CM, (1992). Methods for research on soil borne phytopathopenic fungi, APS Presss, Paul, Minnesota, USA. P. 265.
- [11] Mamatha, T. and Ravishankar, R. V. 2004. Evaluation of fungicides and plant extracts against *Fusarium solani* leaf blight in *Terminalia catappa*. Journal of Mycology and Plant Pathology. 34(2): 306-307.
- [12] Joseph, B., Muzafar, A. D. and Kumar, V. 2008. Bioefficacy of plant extracts to control *Fusarium solani* f. sp. *melongenae* incitant of brinjal wilt. Global Journal of Biotechnology & Biochemistry (2): 56-59.
- [13] Mallesh S. B., Narendrappa, T. and Kumari. 2009. Management of root rot of sage (*Salvia officinallis*) caused by *Fusarium solani* and <u>Rhizoctonia solani</u>. International Journal of Plant Protection. 2(2): 261-264.
- [14] Obongoya, B. O., Wagai, S. O. and Odhiambo, G. 2010. Phytotoxic effect of selected crude plant extracts on soil-borne Fungi of common bean, *African Journal of Crop Science*. **18**(1): 15-22.
- [15] Patel, N. N., Joshi, K. R., Patel, P. M., Patel, M. R. and Patel, M. R. 2010. Bio-efficacy of plant extracts against *Fusarium solani*. *International Journal of Plant Protection.* **3** (2): 306-308.
- [16] Preethi F, Fernandes J, Pricilla K. 2010. Hypoglycemic activity of *Saraca indica* Linn. Bark. J Pharm Res. 3:491-493.
- [17] Maruthappan V, Shree KS. 2010. Antiulcer activity of aqueous suspension of *Saraca indica* flower against gastric ulcers in albino rats. J Pharm Res. 3:17-20.
- [18] Sharma A, Gupta S,Sachan S,Mishra A, Banarji A. 2011. Anthelmintic activity of the leaf of *Saraca indica* Linn. Asian J Pharm Life Sci. 1:391-395.

- [19] Sainath RS, Prathiba J, Malathi R. 2009. Antimicrobial properties of stem bark of *Saraca indica*. Eur Rev Med Pharmacol Sci. 13:371-374.
- [20] Hawas UW, Eltomy SA, AbouZid S, EI-Hossary GA, Nassif RM. 2012.Lipid content and antimicrobial activity of some Egyptian Fabaceae (Leguminoceae) plants. J Med Plants Res. 6(44):5604-5608.
- [21] U. P. Singh, H. B. Singh, R. B. Singh. The fungicidal effect of neem(*Azadirachta indica*) extracts on some soil borne pathogens of gram(*Cicer arietinum*). Mycologia/ Vol. 72, No. 6:1077-1093; Nov-Dec 1980.
- [22] H. N. P. Singh, M. M. Prasad, K. K. Sinha. Efficacy of leaf extracts of some medicinal plants against disease development in banana. Letter in Applied Microbiology, 17(1993), PP. 269-271.
- [23] Kashyapa K, Chand R. 2006. The useful plants of India. CSIR, New Delhi: National Institute of Science Communication and Information Resources.
- [24] Yaye YG, Ackah JAAB, Kra AKM, Djaman AJ (2012). Antifungal activity of different extracts of Terminalia mantaly H. Perrier on the in vitro growth of *Aspergillus fumigatus*. European Journal of Sciences Resources 82: 132-138.
- [25] Quattara S, Kporou KE, Kra AKM, Zirihi GN, N' guessan JD, coulibaly A, Djaman AJ (2013a) Antifungal activities of Terminalia ivorensis A. Chev. bark extracts against *Candida albicans* and Aspergillus fumigatus, Journal of Intercultural Ethnopharmacology 2:49-52.
- [26] Kra AKM, Ahon GM, Djo-Bi D, Ouattara s, Coulibaly A, Djaman AJ (2014) Antifungalactivities of medical plants extracts of Ivorian pharmacopoeia. Journal of Intercultural Ethnopharmacology3:159-166.
- [27]Ellis, M.B (1971). Dermatiaceous Hyphomycetes. Commonweaith Mycological, Kew, Surrey, UK.
- [28]Ellis, M.B (1976). More Dermatiaceous Hyphomycetes. Commonwealth.
- [29] Domsch, K. H., Gams, W and Anderson, T. H. (1993). Compendium of Soil Fungi. IHW-Verlag, Eching, Germany.
- [30] Cheesbrough, M. (2000). District Laboratory Practice in Tropical Countries Part 2, Cambridge University Press, Cambridge, P. 47-54.
- [31] Abbey, S.D. (2007). Foundation in Medical Mycology, 4th edn Kenalf Publication, Port Harcourt, Nigeria, p 22-30.
- [32] Tuite, J. Plant Pathological methods: Fungi and bacteria. Minnesota. Burgess Publishing 1969.
- [33] Sinha K. K., Sinha A. K., Prasad G., 1993. The effect of clove and cinnamon oils on the growth and aflatonin productions by Aspergillus flavus. Letters in Applied Microbiology 16: 114-117.
- [34] Singh J, Tripathi N. N. 1999. Inhibition of storage fungi of blackgram (*Vigna mungo*) by some essential oils. Flavour and fragrance Journal, 14:1-4.

Author Profile



Pushpa Kumari, B.Sc (Botany), MSc. (Botany), Patna University. Research scholar pursuing Ph.D under the able guidance of Reena Mohanka from Patna University.

Volume 7 Issue 7, July 2018

www.ijsr.net

Licensed Under Creative Commons Attribution CC BY

International Journal of Science and Research (IJSR) ISSN (Online): 2319-7064 Index Copernicus Value (2016): 79.57 | Impact Factor (2017): 7.296



Priyanka, B.Sc (Botany), MSc. (Botany), Patna University. Research scholar pursuing Ph.D under the able guidance of Reena Mohanka from Patna University.

Dr. Reena Mohanka, Associate Professor, Science College, Patna University.