Study on the Pattern of Rainfall Distribution using Some Competing Probability Distribution in Cooch Behar District of West Bengal

Rakesh Kumar Meena¹, Sharda Dubey², S. Basak³

1. Introduction

The agriculture land in our country amounts to 143.8 million hectare out of which 67% is dry land and rain fed. The dry land agriculture contributes 44% of our national food grain production. In West Bengal, nearly 72% of the population lives in rural areas. Agriculture is the predominant occupation in the state. The total reporting area of the state is 86.84 lakh ha. of which 52.96 lakh ha that is nearly 61% is the net sown area. The grossed cropped area is 97.52 lakh ha with a cropping intensity of 84%. The climate of the state is tropical and humid except in northern Himalayans.

The climate of the Cooch Behar district of west Bengal, in general, is sub-tropical per humid to tropical humid in nature with distinctive characteristics of high rainfall, high relative humidity and low temperature. About 70% of the annual rainfall is received during the southwest monsoon season, June being the rainiest month. On an average there are about 102 rainy days with records of more than 400 mm rainfall in 24 hours. During monsoons, starting from June, Cooch Behar experiences heavy rainfall. The average annual rainfall is about 3201mm in Cooch Behar district of West Bengal. The numbers of rainy days in Cooch Behar district is recorded to be the highest in the months of June and July. In Cooch Behar district, the number of rainy days often crossed 20 in a month during the months of June and July followed by 15-20 in August and September. Which during April, May and October is varies maintaining a central tendency towards 10 in each month. Also the above mentioned information is based on rainfall data of 1971-1981. An early onset of pre- monsoon shower and occurrence of a low quantum of precipitation during the month of March and April with occasional short dry spells are the other distinctive characteristics for the district of Cooch Behar.

The primary objectives of this research are to estimate and evaluate distribution parameters that may be used to describe the probability of monthly rainfall accumulation for a More specifically, probability location. distribution parameters are estimated from monthly model derived historical rainfall values with a spatial resolution compatible with current agro climatic models, and the goodness-of-fit of the parameters assessed. Direct interpretation of these parameter estimates results in a broad technique for the description of general rainfall regimes for the entire continent. The result of this research is a new portrayal of African rainfall utilizing these probability distribution parameters. In addition to the interpretation of these parameters, description of potential uses of the parameters in hydrologic resource modeling is introduced. Probabilistic information can be utilized to dynamically evaluate rainfall accumulations as well as test different scenarios that can be used as input into other models.

2. Material and methods

Daily rainfall data were collected from Gramin Krishi Mausam Seva Project, AMFU-Pundbari, Cooch Behar for the rain gauge station at AMFU-Pundbari from the year 1995 to 2015 (20 years). Rainfall was measured with IMD specified manual and automatic rain gauge.

To study the characteristics of the rainfall the individual year will be grouped according to the weeks of commencement of rainy season. The data will then, be processed to identify the maximum rainfall received on any one day (24 hours duration), in any week (7days), in a month (4 weeks), in a monsoon season (4 months), in a year (365 days period). Also we calculate total rainfall for all years, the mean, mode and median date, standard deviation and coefficient of variation regarding onset withdrawal and duration of south west monsoon.

Methods of estimating rainfall probabilities:

In estimating probability, two different methods may be tried. Viz.,

- 1) Past frequencies are directly taken to delimit future expectations and
- 2) A theoretical distribution may be fitted to available data and probable frequency may be found out from such a distribution.

Frequency distribution of weekly rainfall is highly skew and includes a number of zero values also. Thom(1966) suggested the fitting of Gamma distribution to data like rainfall having zero lower bound. Mooley (1973) studied Asian summer monsoon rainfall by fitting Gamma distribution on monthly rainfall total. In the present study rainfall probabilities are computed by fitting mixed Gamma distribution to weekly totals. It is given by: G(X) = q + pF(X) where F(X) is the Gamma distribution function, q is the probability of zero precipitation and p=1-q.The distribution function F(X) of the two-parameter Gamma distribution is: $F(X) = \int_0^\infty \frac{x^{\alpha-1} e^{-(\frac{x}{\beta})}}{\beta^{\alpha} \Gamma \alpha} \alpha \& \beta$ are shape and scale parameters respectively of the distribution and $\Gamma(\alpha)$ is the Gamma function of α . The distribution is bounded at the left side by zero. G(X) is the cumulative probabilityofrain<X.To obtains the rainfall probabilities the two parameters α and β of the Gamma distribution have to be estimated from the

Licensed Under Creative Commons Attribution CC BY

observed data. Thom (1958) has shown that the maximum likelihood estimates are efficient consistent and jointly sufficient.Wehave used this method.

The following equation has been solved on the computer by iteration process for X when $P_{z} = 0.10, 0.20, \dots 0.90$ respectively utilising the procedure given in detail by Mooley(1973), where, $Z = x/\beta$

$$P_{Z} = 1 - \left(q + p \int_{0}^{z} \frac{z^{\widehat{\alpha} - 1} e^{-z}}{\sqrt{\widehat{\alpha}}} dz\right)$$

The amounts calculated at 10% to 90% probability levels are designated as assumed rainfall(AR) here onwards. For each station, graphs have been prepared showing for each week, the AR values for the nine probability levels, viz., 10%, 20%,.....90%.An examination of these figures for different stations shows that there are one or more high peaks in AR. On the basis of the amplitudes of the maxima and the positions of the peaks in the rainfall march of time, the patterns are classified. Similar rainfall patterns can be grouped together to form homogenous rainfall pattern zones.

Potential evapotranspiration (PET) was that calculated from U.S. class 'A' pan evaporimetervalues multiplied by 0.7 (pan coefficient) (Banik, 1996).

Moisture availability index (MAI) has been calculated and it is defined as the ratio of weekly rainfall at different confidence levels to PET of the corresponding period (Hargreaves, 1971). The MAI was, however, worked out for 30, 50 and 70 percent probability levels. Depending on the threshold values of MAI the agricultural operations for rice cultivation can be recommended (Banik, 1996).

Threshold values of MAI	Agricultural operations
< 0.3	No. agricultural operations can be done
0.3-0.5	Land preparation
0.5-0.75	Direct sowing of rice can be done
0.76-1	transplanting
>1	Without any risk

Gamma Distribution

The time taken for a number β of events to occur in a Poisson process is described by the gamma distribution, which is the distribution of a sum of β independent and identical exponentially distributed random variables. The gamma distribution has a smoothly varying form like the typical probability density function and is useful for describing skewed hydrologic variables without the need for log transformation. It has been applied to describe the distribution of depth of precipitation in storms. The p.d.f. of one parameter gamma distribution is given by f(x) = $\frac{1}{\Gamma(\beta)}u^{\beta-1}e^{-u}$, x ≥ 0 , $\beta \geq 0$. The gamma distribution involves the gamma function $\Gamma(\beta)$, which is given by $\Gamma(\beta) = (\beta - 1)! = (\beta - 1) (\beta - 2)....2.1$. $\Gamma(\beta) = \int_0^\infty u^{\beta - 1} e^{-u} du$. The two-parameter gamma distribution (parameters β and λ) has a lower bound at zero, which is a disadvantage for application to meteorological variables that has a lower bound larger than zero. Maximum intensity values used in IDF analyses have a lower bound larger than zero and hence the gamma distribution has a limitation in IDF analyses.

M.L.E estimation of Gamma: The equations used for computation of minimum assured rainfall are mixed continuous distributions. As for example for Gamma distribution; G(X) = q + p F(X) where F(X) is the Gamma distribution function and q is the probability of zero precipitation and p=1-q. The distribution function F(X) of the two parameter Gamma Distribution is:

$$F(X) = \int_0^x \frac{x^{\alpha-1}e^{-x/\beta}}{\beta^{\alpha\Gamma}\alpha} dx \text{ where } x, \alpha, \beta > 0$$

= 0 when $x \le 0$

 α , β are shape and scale parameters respectively of the distribution. $\Gamma(\alpha)$ is the Gamma Function of α $=\int_{0}^{\infty} x^{\alpha-1} e^{-x} dx$. In the above notation G(X) is the probability of rain < X.

$$\mathbf{L} = -\mathbf{n}.\boldsymbol{\alpha}.\log\boldsymbol{\beta} - \mathbf{n}\log\Gamma(\boldsymbol{\alpha}) + (\boldsymbol{\alpha} - 1)\sum \log x - \frac{1}{\beta}\sum x$$

Where the summation is over the n sample values. Differentiating as indicated in equations, we the find M.L. equations

 $\log \hat{\beta} + \frac{\partial}{\partial \hat{\alpha}} \log \Gamma \hat{\alpha} - \frac{1}{n} \sum \log x = 0$ Since $\frac{\partial}{\partial \hat{\alpha}} \log \Gamma \hat{\alpha}$ is the digamma function, $\psi(\hat{\alpha})$, We may write in the simplified form

Taking logarithms of (1) and substituting for log $\hat{\beta}$ in (2) gives

$$\log \hat{\alpha} - \psi(\hat{\alpha}) = \log \bar{x} - \frac{1}{n} \sum \log x$$

This equation is implicit in $\hat{\alpha}$ but may be solved with some difficulty using the Davis tables of the ψ – functions. Masuyama and Kuroiwa prepared tables of logarithms and tables of the digamma functions.

Theory of OEM &EEM: Raman (1974) has defined the terms 'OEM' and 'EEM' to work out some meteorological parameters in the case of West Bengal state. But the terai zone of West Bengal being pre -dominantly high rainfall (on the average more than 60mm/week) area, the second criterion of Raman's definition has been modified accordingly to suit the situation of two north Bengal districts under consideration. The modified definitions of OEM and EEM as used in used in the present article are as follows: Suppose during the monsoon months of a year, a particular region having average evaporation of 'e' mm, gets 'x 'mm rainfall at the first day during the commencement of seven days' spell with total rainfall of 'x' mm. Then, this seven days' spell satisfying the following three criteria may be defined as the OEM week :

(i)
$$x \ge e$$
,

(ii) $X \ge (7e+10)$ and

(iii) At least four of the seven days have daily rainfall ≥ 2.5 mm.

Similarly, if the region receives x mm rainfall on the last during an end of seven days' spell may be defined as the EEM week, provided the following criteria are satisfied:

```
(i) x' \ge e,
```

```
(ii) X'\geq(7e+10) and
```

(iii) At least four of the seven days have daily rainfall ≥ 2.5 mm.

Volume 7 Issue 8, August 2018

International Journal of Science and Research (IJSR) ISSN (Online): 2319-7064 Index Copernicus Value (2016): 79.57 | Impact Factor (2017): 7.296

According to WMO, the WMO has divided calendar year into12 duration of 30 days each- called periods and 52 weeks –called standard meteorological weeks of 7 days each. The first week starts on 1^{st} January and ends on 7^{th} January and so on. The 9^{th} (26th Feb to 4^{th} march) week in a leap year and the 52^{nd} week consist of 8 days.

3. Results and Discussion

Descriptive statistics: The graph below demonstrates the total annual rainfall at Cooch Behar district over the study

period. Trends observed indicate total rainfall between 4472.6 mm. to 1529.5 mm. per year, a yearly rainfall variation of 23.92 % over the years. Based on 21 years average the annual rainfall is 3200 mm. The year 1998 was the wettest year with 4472.6 mm of rain and the year 2010 was the driest year with 1529.5 mm of rain. A decrease in annual rainfall of 84.54 mm per year is observed. Based on fitted trend line a decrease in yearly total rainfall of 1.79% per year is also seen.



The graph below demonstrates the yearly total wet day's frequency at Cooch-Behar district. An average annual wet day is 107 days which is 29.31% of total number of days in a year. Yearly wet days counts were observed between 117 days to 77 days, a variance of 29 %. The year 1998 records the highest no. of wet days (117 days) and the Year 2010 records the lowest no. of wet days (77 days), a difference of

34% two years apart. The wet day pattern follows a very similar distribution to total rainfall pattern except in the year 2011 & 2012. A sharp increase in total annual rainfall is observed from 2011 to 2012 whereas total number of wet days remains almost same during the same period.



The graph below presents the average monthly rainfall between 1995 to 2015. The observed trend, gradual and consistent increase from January to April followed by a sudden & sharp increase in May & June. It peaks in July and

thereafter a slight drop in August. Thereafter there is a sudden and sharp drop in rainfall in the months of September & October followed by nearly no rain in the months of November & December.

Volume 7 Issue 8, August 2018 <u>www.ijsr.net</u> <u>Licensed Under Creative Commons Attribution CC BY</u>



The month of July records the highest monthly average rainfall 743.48 mm or 22% of total annual rainfall. Whereas the month of November records the lowest monthly average rainfall 6.45 mm or 0.1% of total annual rain closely followed by December at 4.25 mm or 0.1% of annual total.

The graph below present the variation in the monthly rainfall over the study period. The variation is 50% or less in the rainfall months of May to September. However in the months of scanty rain i.e. October to April the variation is more indicating a erratic rainfall months.



The following graph present the monthly average wet days over the study period. The observed trend is same as in case of monthly average rainfall. June, July & August together record the highest no. of wet days (20 each approx.) i.e. 52.33% of annual total.

Wet days of more than 14 days per month occurs 25 times, 8 times each in July & August, 7 times in June & 4 times in May.





Occurrence of rainy days is more uniform from May to September when variation is less than 28%. Occurrence of wet day is more erratic in other months & this is highest in the month of December followed by November & January. The variation is beyond 96%.

Volume 7 Issue 8, August 2018 www.ijsr.net Licensed Under Creative Commons Attribution CC BY



The above mentioned Seasonality Index has been computed over the study period.

Calculating the Seasonal index is giving the formula by Walsh and Lawer (1981):

 $S.I._{k} = \frac{1}{R_{k}} \sum_{n=1}^{12} \left| X_{nk} - \frac{R_{k}}{12} \right|$

Where, X_{nk} = Rainfall of month n of the year k. R_k = total annual rainfall for the year k.

Low value of seasonality index indicates a shorter dry season & a better distribution of monthly rainfall among the months of a year. A high value of S.I. indicates that most of the rain occurs within few months (2-3 months) & a longer (frequent) dry season. Trend analysis has shown 2.5% increase in S.I. per year. An increasing trend is an indicator of alarming situation in Agriculture.

Table 1: Weekly maximum & minimum rainfall year wise

Year	Week No	Maximum	Week	Minimum
		rainfall (mm)	No	rainfall(mm)
1995	33	559.7	10	0.4
1996	28	372.4	13	0.4
1997	24	468.3	6	0.6
1998	36	632.7	9	0.2
1999	28	694.6	50	0.5
2000	27	543.8	44	0.8
2001	34	370.6	16	0.3
2002	29	374.7	4	0.2
2003	26	296.1	1	0.2
2004	29	399.5	42	1.8
2005	25	476.4	47	0.4
2006	27	394.2	43	0.2
2007	36	402.4	11	0.1
2008	35	641.4	6	1
2009	26	283.1	12	0.1
2010	30	134.5	7	0.6
2011	38	344.7	3	0.5
2012	28	371.1	4	0.2
2013	28	381.1	12	1.9
2014	23	376.7	50	0.4
2015	35	302.7	5	0.1

Table 2: I	Date wise &	week wise	maximum	rainfall for
different	years along	with month	ly maximu	m rainfall.

Year	Week	Date of	Maximum	Name of	Monthly
	No	Rainfall	rainfall	Month	Maximum
			(mm)		rainfall(mm)
1995	33	15.08.1995	183.9	August	1061.9
1996	28	13.07.1996	101.8	July	1193.2
1997	24	09.7.1997	136	August	1059.2

1998	36	3.9.1998	284.9	August	948.9
1999	28	10.7.1999	474	July	1274.4
2000	27	06.07.2000	196.3	June	981.1
2001	34	22.08.2001	132.3	September	760.5
2002	29	20.07.2002	200	July	1107.2
2003	26	26.07.2003	96.2	June	935.9
2004	29	11.07.2004	86.6	July	1190.9
2005	25	18.07.2005	53	August	778.5
2006	27	28.07.2006	160	September	506.5
2007	36	30.08. 2007	47.8	July	618.3
2008	35	30.08.2008	239.6	August	1172.7
2009	26	23.06.2009	116.8	June	540.3
2010	30	23.07.2010	39.6	August	360.2
2011	38	18.9. 2011	214.60	July	800.4
2012	28	11.7.2012	65.60	July	739.8
2013	28	10.7. 2013	203.20	July	682.7
2014	23	8.6. 2014	128.90	June	604.3
2015	35	02.09.2015	88.90	August	797.3

 Table 3: Yearly mean dates of OEM & EEM and duration of CDS at Cooch Behar

			102540	0000	Bena	
Year	Date	Week	Date of	Week	Duration of	Week No.
	of	NO.	end	No.	CDS	
	onset					
1995	28	22	23	38	1-10 May	17,18
	May		September		-	
1996	28	22	7	36	18-27	38,39,41,42,43
	May		September		September ;	
					8-17	
					October;18-	
					27 October;	
1997	8	23	20	38	18-27	42,43
	June		September		October	
1998	31	22	5	36	8-17	36,37,38,41,42
	May		September		September	
					;8-17	
					October	
1999	-		-		21-30	42,43,44
					October	
2000	28	22	23	38	18-27	42,43
	May		September		October	
2001	3	22	8	36	18-27	42,43
	June		September		October	
2002	2	22	28	39	18-27	42,43
	June		September		October	
2003	8	23	13	37	-	
	June		September			
2004	13	24	25	39	18-27	42,43
	June		September		October	
2005	12	24	-		8-17 October	41,42
	June					
2006	11	24	23	38	-	
	June		September			

Volume 7 Issue 8, August 2018 <u>www.ijsr.n</u>et

International Journal of Science and Research (IJSR)				
ISSN (Online): 2319-7064				
Index Copernicus Value (2016): 79.57 Impact Factor (2017): 7.296				

2007	3	22	8	36	18-27	42,43
	June		September		October	
2008	2	22	28	39	8-17 October	41,42
	June		September			
2009	1	22	-		18-27	42,43
	June				October	
2010	14	24	6	36	18-27	42,43
	June		September		October	
2011	4	23	23	38	8-17	41,42,43
	June		September		October;18-	
					27October	
2012	3	22	22	38	19-28	42,43
	June		September		October	
2013	4	23	9	36	17-26	38,39,42,43
	June		September		September	
					;17-26	
					October	
2014	4	23	-		29 Sep -8	39,40,41,42,43
	June				October; 19-	
					28 October	
2015	28	22	2	35	8-17	41,42,43
	May		September		October;18-	
					27October	

 Table 4: Estimated parameters of rainfall Characteristics of Cooch Behar

Particulars	Cooch -Behar
Mean date of OEM	4 June
Median date of OEM	2 June
Mean deviation from mean (OEM dates)	4 days
Quatrile deviation (OEM dates)	4 days
Earliest probable date of OEM (p=0.50)	31 May
Latest probable date of OEM (p=0.50)	8 June
Mean date of EEM	17September
Median date of EEM	14 September
Mean deviation from mean (EEM dates)	9 days
Quatrile deviation (EEM dates)	9 days
Earliest probable date of EEM (p=0.50)	8 September
Latest probable date of EEM (p=0.50)	26 September
Average number of dry spell/year (4	8 days
days or more)	
Average deviation of dry days /year	8 days
Average number of CDS /year	1.78
Average amount of rainfall in OEM week	119.09 mm per week
Range (rainfall during OEM weeks)	9.6 – 376.7 mm per week
Average amount of rainfall in EEM week	60.50 mm per week
Range (rainfall during EEM weeks)	0.4 – 632.7 mm per week

Analysis of 21 years' rainfall data, shows OEM between 28th May to 14th June with an average rainfall of 119 mm/week. The mean and median dates of OEM are 4th and 2nd June respectively. The mean deviation from mean and quartile deviation of mean dates of different years are same i.e. 4 days.Hence the earliest and the latest probable dates of OEM are 31st May and 8th June respectively. The monsoon effectively ceased between 2nd September to 28^{th} September with an average rainfall of 60.50 mm/week. The range of amount of rainfall during OEM week in different years is between 9.6 - 376.7 mm per week and that of EEM week between 0.4 - 632.7 mm per week. The total number of CDS is 26 during the years 1995-2015, of which 15 occur during end of October, 6 in mid October, 2 in end of September and

1 each in first part of May, mid of September and 1st part of October.

 Table 5: Monsoon onset (week), withdrawal (week) and length (week) of Cooch Behar

Year	Monsoon	Monsoon	Annual	Length
	onset	withdrawal	rainfall	-
	Week	Week	(mm)	
1995	22	38	4331	16
1996	22	36	2621.1	14
1997	23	38	3848.7	15
1998	22	36	4472.6	14
1999	-	-	4230.6	-
2000	22	38	3269.3	16
2001	22	36	3615.4	14
2002	22	39	2978.5	17
2003	23	37	3486.8	14
2004	24	39	3521.7	15
2005	24	-	3710	-
2006	24	38	2571.6	14
2007	22	36	2808.7	14
2008	22	39	3941.8	17
2009	22	-	2739.2	-
2010	24	36	1529.5	12
2011	23	38	2853.5	15
2012	22	38	3422.3	16
2013	23	36	2451.5	13
2014	23	-	2115.5	-
2015	22	35	2640.12	13
Mean	22.65	37.23	3198.068	14.65
Mode	22	38	-	14
S.E.±	0.813	1.3	765.16	1.41

The average onset of monsoon is 23rd SMW (4 -10th June). Its range varies from 22nd - 24th SMW over the study period. Monsoon commenced as early as 22nd SMW in1995, 1996, 2000 (28th May), 1998 (31st May), 2009 (1st June), 2002,2008 (2nd June), 2001, 2007 (3rd June). Late onset of monsoon is found on 24th SMW in 2006 (11th June), 2005 (12th June), 2004 (13th June) & 2010 (14th June). The average withdrawal of monsoon is in the 37th SMW (10 -16th September). The range of withdrawal of monsoon is 35th SMW to 39th SMW. The earliest withdrawal of monsoon is observed on 2nd September in 2015 and it is as late as 28th September in 2002, 2008& 25th September in 2004.

Finding of different distributions: The probability distributions viz. normal, lognormal, gamma, weibull, Exponential was identified to evaluate the best fit probability distribution for rainfall. In addition the different forms of these distributions were also tried and thus the probability distributions viz. normal, lognormal, gamma, weibull, Exponential were applied to find out the best fit probability distribution.

Identifying the best fit distribution: The three goodness of fit test mentioned above were fitted to the maximum rainfall data treating different data set. The test statistic of each test were computed and tested at (α =0.01) level of significance. Accordingly the ranking of different probability distributions were marked from 1 to 16 based on minimum test statistic value. The distribution holding the first rank was selected for all the three tests independently. The assessments of all the probability distribution were made on the bases of total test

Volume 7 Issue 8, August 2018 www.ijsr.net

score obtained by combining the entire three tests. The total score of the entire three tests were summarized to identify the best fit distribution on the bases of lowest score obtained.

Finally the best fit probability distributions for maximum rainfall on different sets of data were obtained and the best fit distribution for each set of data was identified. The above methodology can also be used for studying the probability distribution pattern for other weather variables.

Table 6: Best Fitted Distribution and Parameter Year-Wise

YEAR	Best fit Distribution	Parameter
1995	Gamma	α=0.48772 β=239.85
1996	Gamma	α=0.59529 β=190.09
1997	Weibull	α=0.53681 β=58.779
1998	Gamma	α=0.74914 β=161.35
1999	Gamma	α=0.8157 β=172.88

2000	Lognormal	σ=1.3791 μ=3.8863
2001	Weibull	α=0.58727 β=83.639
2002	Gamma	α=0.66309 β=121.4
2003	Weibull	α=0.56535 β=72.473
2004	Weibull	α=0.76297 β=85.969
2005	Gamma	α=0.78037 β=120.8
2006	Gamma	α=0.52804 β=131.62
2007	Gamma	α=0.67581 β=118.75
2008	Gamma	α=0.71235 β=153.71
2009	Gamma	α=0.96824 β=94.467
2010	Weibull	α=0.66646 β=46.101
2011	Weibull	α=0.64819 β=71.71
2012	Weibull	α=0.62111 β=83.589
2013	Weibull	α=0.74898 β=58.599
2014	Gamma	α=0.69251 β=98.959
2015	Gamma	α=0.90729 β=91.806



Figure 1: Density graph of Gamma distribution

 Table 7: Goodness of Fit (study period 1995-2015)

 Summary

		Kolmogorov		Anderson		Chi-Squared	
#	Distribution	Smirnov		Darling			
		Statistic	Rank	Statistic	Rank	Statistic	Rank
1	Exponential	0.12124	3	30.843	4	126.06	4
2	Gamma	0.03843	1	2.1812	1	34.624	2
3	Lognormal	0.13382	4	18.71	3	110.9	3
4	Normal	0.20793	5	52.63	5	154.5	5
5	Weibull	0.06559	2	3.0505	2	28.505	1

 Table 8: Fitting Results (study period 1995-2015)

#	Distribution	Parameters
1	Exponential	λ=0.01074
2	Gamma	α34366.0= β92.041=
3	Lognormal	σ 1958.1 = μ 4425.3=
4	Normal	σ 72.411 = μ 270.39=
5	Weibull	α64476.0= β 101.97=

Table 9: Number of years of monsoon	onset and	withdrawal
at Cooch Behar		

ut Coolen Denta							
Onset of monsoon	Withdrawal of Monsoon						
	Early	Normal	Late				
	(2-10 Sept.)	(11-19 Sept.)	(20-28 Sept.)				
Early (28-2 June)	3(4249.60)	0	4 (3750.43)				
Normal(3-8 June)	3(2958.53)	1(3486.8)	3 (3635.5)				
Late(9-14 June)	1(1529.5)	0	2(3046.65)				
Amount of rainfall (mm)	2912.54	3486.8	3477.52				

*Figure within the parenthesis is the average annual rainfall in mm.

The total annual rainfall is more than average (3198 mm) during the year in which monsoon onset is observed in 22nd and 23rd SMW. It is less than average (1529 -3046 mm) when onset is around 24th SMW. Heavy deficit in rainfall (1529.5 mm) is found when commencement of monsoon is late (around 24th SMW)but withdrawal of monsoon is early (36th SMW). Total annual rainfall is highest (4249.6 mm)when there is early commencement (22nd SMW) as well as early withdrawal (36th SMW) of monsoon. The

Volume 7 Issue 8, August 2018

www.ijsr.net

second highest annual rainfall (3750 mm) is observed when onset of monsoon is early but its withdrawal is late. With early to normal onset of monsoon and early withdrawal the average number of dry spell is more. The earliest commencement of dry spell is observed in 1995 when there is late withdrawal of monsoon. However most CDS occur late and its occurrence varies from 36th to 44th SMW.



Figure 2: Occurrence of dry spells (≥4 days)

Table 10: Weekly best fit distribution	& its parameter
estimates	

Week no	Best fit dist	Parameter estimate
10	Log- logistic	α=0.82435 β=0.70477
11	Log normal	σ=1.8401 μ=1.3782
12	Log normal	σ=1.9756 μ=1.5481
13	weibull	α=0.64761 β=31.478
14	Log- logistic	α=1.9269 β=26.758
15	Gamma	α=1.3114 β=17.587
16	Gamma	α=1.9469 β=21.41
17	Normal	σ=35.96 μ=47.853
18	weibull	α=0.87393 β=61.01
19	Gamma	α=1.7363 β=30.057
20	Normal	σ=47.487 μ=57.9
21	Log normal	σ=0.77245 μ=4.3988
22	Normal	σ=77.307 μ=127.54
23	weibull	α=1.2027 β=142.82
24	Exponential	λ=0.00681
25	weibull	α=1.3152 β=225.7
26	Log normal	σ=0.76464 μ=4.7152
27	Gamma	α=1.7926 β=109.41
28	Exponential	λ=0.00465
29	Gamma	α=0.86931 β=168.96
30	Exponential	λ=0.0081
31	Gamma	α=1.3447 β=69.256
32	weibull	α=1.0134 β=110.85
33	Exponential	λ=0.00502
34	Log normal	σ=1.0367 μ=4.5407
35	Log- logistic	α=1.8568 β=83.408
36	weibull	α=0.82011 β=116.61
37	Gamma	α=0.98831 β=88.867
38	Gamma	α=0.79016 β=142.86
39	Gamma	α=1.9844 β=51.608
40	Exponential	λ=0.01568
41	Exponential	λ=0.01422
42	Log- logistic	α=0.86575 β=11.066

Table 11: Long term average weekly rainfall (mm) at
different probability levels of Cooch Behar

different probability levels of Cooch Benar								
week	Average	P=0.3	P=0.4	P=0.5	P=0.6	P=0.7		
rainfall (mm)								
10	0.77	1	0	0	0	0		
11	5.30	2	0	0	0	0		
12	11.35	7.6	4	1.8	0.3	0		
13	28.65	25	13.5	6.35	2.1	0		
14	26.30	25.7	19	12.85	5.1	0		
15	20.86	26.05	20.1	15.35	11.3	7.7		
16	35.7	46.1	37.05	29.4	22.55	15.9		
17	43.3	56.8	42.9	29.6	15.8	0		
18	60.3	71.8	51.8	36.8	25.1	15.8		
19	49.8	61.7	50	40.4	31.95	24.2		
20	57.9	77.8	62.1	47.4	32.8	17		
21	108.1	21.5	6.55	2.15	0	0		
22	127.6	144	109.6	77	44.8	10.4		
23	138.7	166.4	132.6	105.2	61.6	60.6		
24	139.9	169.4	127	94.4	67.6	4.5		
25	205.1	259.4	210.8	170.6	135.4	102.9		
26	145.6	25.2	7.05	2.13	0	0		
27	196.1	239.4	196.4	161	130.2	102.2		
28	215.2	258.4	196.8	148.8	109.6	76.4		
29	146.9	174.3	129.8	95.6	68.4	46		
30	117.1	148.4	113	85.5	63	44		
31	93.1	113.4	90	71.3	55.4	41.3		
32	110.6	132.8	101.6	77	57	40		
33	199.1	239.4	182.4	138	101.6	70.8		
34	153.0	30.2	8.85	2.8	0.89	0		
35	137.8	0	0	5.8	0	0		
36	139.3	146	104.8	74.4	51.4	33.1		
37	87.8	105.6	80.2	60.5	44.5	31		
38	107.5	126.3	91	64.4	43.5	26.8		
39	97.5	121.2	99.6	81.6	65.8	51		
40	54.7	67	48.6	34.4	22.7	12.9		
41	53.6	45.2	45.3	29.6	16.8	5.9		
42	29.9	9.6	0.145	0	0	0		

Rainfall at different confidence levels show that rain started around10th SMW at 0.3 confidences level whereas it is in the 15th SMW when confidence level is 0.7. The highest PET is 24.78 mm/week in 17th SMW i.e. in the last week of May when monsoon is about to arrive. The higher values of PET (i.e. more than 14) are observed from 11th SMW i.e. middle of March to 41st SMW i.e. 2nd week of October when late autumn season is about to arrive. Once late autumn begins the PET values sharply decrease. The lowest is 3.745 mm/week at 2nd SMW i.e. in the 2nd week of January which is at the middle of winter season.

International Journal of Science and Research (IJSR) ISSN (Online): 2319-7064 Index Copernicus Value (2016): 79.57 | Impact Factor (2017): 7.296



Figure 3: Plot of MAI values vs. week no. at different probability levels.

Table 12: Long term Moisture Availability index at different probability levels of Cooch Benar								
Week	Average MAI	MAI P=0.3	P = 0.4	P=0.5	P=0.6	P = 0.7		
10	0.0635	0.082576	0	0	0	0		
11	0.298	0.112486	0	0	0	0		
12	0.717	0.480405	0.252845	0.11378	0.018963	0		
13	1.912	1.668892	0.901202	0.423899	0.140187	0		
14	1.692	1.653797	1.222651	0.826898	0.328185	0		
15	1.33	1.661352	1.281888	0.978954	0.720663	0.491071		
16	1.783216783	2.302697	1.850649	1.468531	1.126374	0.794206		
17	1.747376917	2.292171	1.731235	1.194512	0.637611	0		
18	3.445714286	4.102857	2.96	2.102857	1.434286	0.902857		
19	3.014527845	3.734867	3.026634	2.445521	1.934019	1.464891		
20	3.308571429	4.445714	3.548571	2.708571	1.874286	0.971429		
21	7.353741497	1.462585	0.445578	0.146259	0	0		
22	7.659063625	8.643457	6.578631	4.621849	2.689076	0.62425		
23	8.540640394	10.24631	8.165025	6.477833	3.793103	3.731527		
24	7.701624002	9.325626	6.991467	5.196807	3.721442	0.247729		
25	13.50230415	17.07702	13.87755	11.23107	8.913759	6.774194		
26	9.904761905	1.714286	0.479592	0.144898	0	0		
27	13.53347136	16.52174	13.55418	11.11111	8.985507	7.05314		
28	13.48370927	16.19048	12.33083	9.323308	6.867168	4.786967		
29	8.799041629	10.44025	7.774783	5.726265	4.097035	2.755316		
30	7.088377724	8.983051	6.840194	5.175545	3.813559	2.663438		
31	5.46201232	6.652977	5.280141	4.183045	3.25022	2.422998		
32	6.929824561	8.320802	6.365915	4.824561	3.571429	2.506266		
33	11.95078031	14.36975	10.94838	8.283313	6.098439	4.2497		
34	10.87420043	2.146411	0.628998	0.199005	0.063255	0		
35	9.156146179	12.69103	8.770764	6.232558	4.451827	3.076412		
36	7.653846154	8.021978	5.758242	4.087912	2.824176	1.818681		
37	5.453416149	6.559006	4.981366	3.757764	2.763975	1.925466		
38	5.952380952	6.993355	5.03876	3.565891	2.408638	1.483942		
39	6.19047619	7.695238	6.32381	5.180952	4.177778	3.238095		
40	3.601053325	4.410797	3.199473	2.264648	1.494404	0.849243		
41	3.496412264	2.948467	2.95499	1.930855	1.09589	0.384866		
42	2.440816327	1.216327	0.644898	0.334694	0.152653	0		

Fable 12: Long term Moisture Availability Index at different probability levels of Cooch Behar

From the above plotted graph for MAI it is observed that land preparation for olitorius Jute and direct seeded Aus rice can be started safely on 18th SMW i.e. around 1st week of May. The MAI values at the 21st SMW (i.e. end of May) are less than 0.3 at probability levels P \geq 0.5. This indicates that no farming operation can be done during that week. Sowing of Aman rice can be started from 22nd SMW onwards as the MAI values are more than 2 at all probability levels except at P=0.7 when MAI is at the lower limit of the range. However its sowing can be safely started from the 23rd SMW onwards. Again there is a sudden drop in moisture level in the 26th SMW when even no farming operation is possible at all probability levels P \geq 0.4. It implies that transplanting of rice cannot be started before 27th SMW. Another drop in moisture level is found in the 34th SMW when even no farming operation is possible at all probability levels $P \ge 0.4$. So the flowering stage of rice (considered to be most critical) should be over before 34th SMW. Again harvesting of jute must be completed by the beginning of 33rd week so that sufficient standing water is available in the canals and other low lying areas for retting of jute. This is possible if farmers can complete sowing of jute by beginning of 16th week strictly as olitorius is a 120 days crop.

Volume 7 Issue 8, August 2018 <u>www.ijsr.net</u>

DOI: 10.21275/ART2019364

4. Summary and Conclusion

For entire period of study although the average onset of monsoon in this area is closer to that of Kerala (normal monsoon date according to IMD, June 2) when the southwest monsoon hits its first spell in the Indian Peninsula, the average retreat of monsoon in this part of Terai zone of West Bengal occurs 2 weeks advance than withdrawal of monsoon rains from the rest of India.During 1995 to 2015 the onset of effective monsoon is delayed by one month as compared to the study period of 1971-1990. On the other hand the withdrawal of monsoon is two weeks advance in comparison to 1971-1990 (Das et al, 1996). Thus the monsoon spell is squeezed by one and half month.It maybe assumed that the arrival of monsoon rains over this part of West Bengal has relation with the onset of monsoon at Kerala. As a result, contrary to other parts of West Bengal, this region experiences about 3.5 months' duration of monsoon rains, i.e., on the average only 2 weeks more than southern part of West Bengal (Das et al, 1996). The observed variation in the OEM date is only 4 days as compared to 9 days for the period of 1971-1990. However the variation in EEM dates for both of the periods is same. In this region, there is possibility of occurrence of dry spell of 10 days or more once in every year, on the average ; the starting date of this dry spell being any time during around 1st and 2nd week of May, i.e. 10 days from the day of onset of monsoon. Again, such a dry spell may appear once in every year, on the average, around 2nd and 3rd week of September. There is also possibility of another such spell of dry days in this area in every in every year, on the average, anytime during 2nd and 4th week of October. The average number of dry spell per year is found to be more in comparison to the study period of 1971-1990. The average amount of rainfall observed in OEM week are almost same for both of the periods but during 1995-2015 it is less than half of the rainfall in EEM week during 1971-1990. The ability of the gamma distribution and parameter estimates to adequately fit the empirical distribution of values in the entire study period have been tested using the KS goodness of fit test. Gamma and Weibull distributions are the best fitted distributions for the rainfall years in Terai region of Cooch Behar district of West Bengal. The ability to represent the rainfall using the gamma distribution parameters allows for interpretation of the parameter estimates as a compact summary of the full rainfall distribution, through analysis of the distribution parameters, it is possible to estimate the likelihood of an area receiving rainfall amounts that would cause flooding, wash out dams or provide sufficient water to support crops.Different agricultural operations can be scheduled using different assured rainfall probabilities. Sowing of jute must be completed by the beginning of 16th week so that sufficient water is available for retting. Sowing of direct seeded Aus paddy can be safely done from 18th SMW onwards. Sowing of Aman paddy can be started from 22nd SMW onwards but its transplanting cannot be done before 27th SMW.Observed critical dry spell duration can be used as an indicator of drought risk. This research could prove valuable to a wide range of groups from scientists studying assessing precipitation, to policy makers forecast information, to farmers in crop planning.

5. Acknowledgement

Mere infinite thanks must not suffice to convey the sense of gratefulness and indebt ness from the core of my heart to the respected chairman of my advisory committee, Dr. S. Basak, Lecturer, Department of Agricultural Statistics, Uttar Banga Krishi Viswavidyalaya, Pundibari, CoochBehar for his auspicious and affectionate supervision, wholesome advice and encouragement, valuable suggestions, and untiring endeavor, creative criticism, abiding interest and mountain sources of inspiration throughout the course of investigation.In thanking all those people, I must emphasis that I am fully responsible for any shortcoming that remains.

References

- [1] AasthaGulati, Palak Gupta, Meenaksh. Jha, P. ParthSarthiand Kumar Vishal,2008. "Impact of climate change, variability, and extreme rainfall elements on agricultural production and food insecurity in orissa."
- [2] Banik, P. and Sharma, R.C. (2009): Rainfall pattern and moisture availability index in relation to rice crop planning in eastern plateau region of India. Journal of Agro meteorology 11(1):54-58
- [3] Barron, J., J. Rockström, F. Gichuki and N. Hatibu (2003). "Dry spell analysis andmaize yields for two semi-arid locations in East Africa." Agricultural and Forest Meteorology 117(1-2): 23-37.
- [4] Bhargava, P.N. The influence of rainfall on crop production .Research Journal of JawaharLalKrishi Vishwavidyalaya.1974; 32, No.1 and 2.
- [5] Das et al.,(1996) "A study on effective monsoon and dry spells of short return periods during monsoon months in two north Bengal districts." Mausam,47,2,PP.145-148.
- [6] Fisher R.A. (1924) The Influence of the rainfall on the yield of wheat at Rothamsted. Philosophical transaction of the royal society of London.1924; series B, vol.123.
- [7] Hazra, A., Bhattacharya, S., Banik, P., (2014)
 "ModellingNakshatra –Wise Rainfall Data of the eastern plateau region of India." MAUSAM, 65, 2 pp.-264-270;
- [8] Kamruzzaman, M., S. Beecham and A. V. Metcalfe (2011)."Non-stationarityinrainfall and temperature in the Murray Darling Basin."HydrologicalProcesses25(10): 1659-1675.
- [9] Maddison, D. (2007). "The perception of and adaptation to climate change in Africa."World.
- [10] Manning, H.L. (1950) Confidence limits of monthly rainfall. Jour, Agril. Sci.1950;40:169.
- [11] Moaley, D.A. (1972) "Gamma distribution probability model for Asian Summer Monsoon monthly rainfall." Monthly weather Review 1972 vol.101, No.2.
- [12] Muralidharan et al.,(2005) "Statistical modelling of rainfall data using modified Weibull distribution."MAUSAM,56,4,PP.765-770.
- [13] Palmer, T. N. and D. Anderson (1994)."The prospects for seasonal forecasting—Areview paper." Quarterly Journal of the Royal Meteorological Society120(518): 755-793
- [14] Palmer, W. C. (1965). Meteorological drought, US Dept. of Commerce, WeatherBureau.

- [15] Rahim, K.M.B., Majumder, D., Biswas, R.k., 2011. "Determinants of Stagnation in Productivity of Important Crops in West Bengal."Study No. 162.
- [16] RanaIsha, Randhawa, S.S., 2010 "Impact of rainfall on agriculture in india".
- [17] Singandhupe, R.B., Anand P.S.B. and Kannan K. (2000).Effects of rainfall pattern on rice productivity in state of Orissa. Crop research, 20(3): 360-366
- [18] Smith, R.L., (1994), "Spatial modelling of rainfall data. In, statistics for the Environment 2: water related issues, edited by V. Barnett and F. Trukman, John Wiley, Chi Chester, 19-42.
- [19] State Agriculture Plan, West Bengal Prepared by NABARD (Specified year 2009-10)
- [20] Suppiah, R. and K. Hennessy (1998). "Trends in total rainfall, heavy rain events andnumber of dry days in Australia, 1910-1990." International Journal of Climatology18(10): 1141-1164.
- [21] Thom, Herbert C. S., "A Note on the Gamma Distribution,"*Monthly Weather Review*, Vol. 86, No. 4, Apr. 1958, pp. 117-122.
- [22] Tsakiris, G., D. Pangalou and H. Vangelis (2007). "Regional drought assessment based on the Reconnaissance Drought Index (RDI)." Water resources management 21(5): 821-833.
- [23] Walsh, R.P.D. and Lawer, D.M.(1981) : Rainfall seasonality : Description, spatial patterns and change through time. Weather, 36, pp.201-208.
- [24] Wilks, D. S., Wilby, R. L. The weather generation game: a review of stochastic weather model *Progress in Physical Geography*, No 23(3), 1999, pp.329–357.
- [25] Wong, G., H. Van Lanen and P. Torfs (2013).
 "Probabilistic analysis of hydrologicaldrought characteristics using meteorological drought." Hydrological SciencesJournal 58(2): 253-270.

DOI: 10.21275/ART2019364

302