

Calculation of Uranium Concentrations in Tikrit University Soil Samples using Neutron Activation and CR-39 Detector

Fareed M. Mohammed¹, Saed S. Kamon², Khalid H. Mahdi³, Daham. M. Nael⁴

^{1,4}University of Tikrit, College of Science, Iraq

²College of madinat al Elem ALJamea, Iraq

³University of Baghdad, College of Education for pure sciences, Ibn al-Haytham, Iraq

Abstract: *The aim of this work was to measure the concentrations of uranium in soil samples collected from the University of Tikrit, which located in the province of Salah al-Din, (180) km north of Baghdad, in this study 80 samples was collected from (14) college in Tikrit University. The samples were preperated with CR-39 for irradiation with thermal neutrons and compared the results with standrad samples, in order to measure uranium concentrations in soil samples. The results show that the highest uranium concentration was (7.31) ppm in sample (VI) college of arts which and the lowest concentration of uranium was (0.54) ppm in the sample (G1) which college Engineering and the weighted average of all samples was (3.48 ± 1.57) ppm, These values were within the universally accepted limit of (11.7) ppm.*

1. Introduction

Human body is usually exposed to radiation with vary rates, whether it is natural or industrial. The effect of the radiation in the environment may remain several years, that change the genetic of humans and animals, leading to genetic mutations that appear in subsequent generations, the impact of contamination should not be neglected in soil and water that transfer from it to human and animal food . [1]

Uranium is a natural component of soil and water. Its concentration ranges approximately from (1_120) ppm and consists of a number of isotopes, ²³⁸U, ²³⁵U, and ²³⁴U. Most of these isotopes emit (α or β) particles and Gamma radiation.[2]

Therefore, there is a need to study the effect of radiation and detection the natural and industrial radiation activity, the extent of pollution of the environment and how to be treated, many researches calculated concentrations of radioactive materials in the soil, plant, building materials, water, air and others, and the indication of their impact on living organisms. [3]

For this purpose, different techniques were used, such as spectral measurements of gamma rays, neutron activation analysis, X-ray fluorescence, and solid state nuclear track detector. [4]

The purpose of this research is to calculate the concentrations of uranium in soil of University of Tikrit (as it attracts an increasing number of students).

2. Experimental Part

a) Collection of Samples

The University of Tikrit is located north of the city of Tikrit in the province of Salah al-Din and within the coordinates listed in Table (1), shown in Figure (1)

Soil samples were collected from 14 college in University of Tikrit at a depth of 5 cm and stored in special bags with 500 g for each sample. Six samples were taken from each college except the printing department only two samples, the distance between the sample and the specific building varies between (2-20)m. This process was carried out at 16/10/2017.




Figure 2: The tape which consists of 25 samples and detector (CR-39)




Figure 3: The irradiation tape which consists of (25) samples with the (CR-39) detector using Neutron source




Figure 4: The track of Nuclear Fission Fragments in the Solid stat Nuclear detector type (CR-39)

e) Measurement of uranium concentrations

The track density was measured by using microscope type (Novel) with a magnification power of (40x) using the following equation [7]:

$$\text{Tracks density } (\rho) = N_{ave}/A \text{ ----- (1)}$$

Where:

ρ : Track intensity (Tracks / mm²)

N_{ave} : Average total number of tracks (Track)

A: Aerea of the field vision (mm²)

Figure (4) shows the track of nuclear fission fragments in the solid state nuclear detector type (CR-39).

The concentration of uranium is measured in soil samples by comparing the density of tracks that recorded by CR-39 for

samples and for standard samples according to the following equation [7] :

$$C_x = C_s(\rho_x/\rho_s) = \rho_x/\text{slope} \text{ ----- (2)}$$

Where:

C_x : Concentration of uranium in the unknown sample (ppm)

C_s : Concentration of uranium in the standard sample (ppm)

ρ_x : Track intensity in the unknown sample (Tracks / mm²)

ρ_s : Track intensity in the standard sample (Tracks / mm²)

The slope of the straight line for the track density and uranium concentration shown in Figure (5) [5].




Figure 5: The relationship between the tracks density and the concentration of uranium (ppm) for standard soil samples.[5]

3. Results and Discussion

For monitoring is necessary to measure the uranium concentration in soil of University of Tikrit , The monitoring at this time and in the future is very important to determined the risks that cause by this concentrations to ensure that its safety .

Table (2) shows the results of uranium concentrations in soil samples at the University of Tikrit. It is clear that the uranium concentration of the study areas ranges between (0.54 and 7.31)ppm. The highest concentration of uranium was (7.31ppm) in samples (V1) which was one of the samples from college of arts and the lowest concentration of uranium found in the samples (G1) from college of Engineering was (0.54) ppm. the weighted average of all samples was (3.48 ± 1.57)ppm, and these values were within the universally accepted limit of (11.7) ppm According to(UNSCEAR 2000) [8]. Figure (6) shows of the results that obtained from this study.

Table 2: Track density and concentration of uranium for all samples of soil and its locations

No.	Locations	Sample cod	Track density (track/mm ²)	Uranium Concentration (ppm)
1	College of Science	S1	2594.2±36.96	6.3
		S2	1924.2±30.57	4.67
		S3	945.7±18.42	2.29

2	College of Education for Sciences of drainage and humanity	S4	2261.4±46.13	5.49
		S5	2142.8±43.11	5.21
		S6	1002.8±14.32	2.43
		T1	1315.7±27.40	3.19
		T2	1591.4±24.75	3.87
		T3	1884.2±36.34	4.58
		T4	1322.8±22.26	3.21
3	College of Engineering	T5	1274.2±89.72	3.09
		T6	421.4±12.30	1.02
		G1	224.2±8.68	0.54
		G2	942.8±22.38	2.29
		G3	548.5±21.70	1.33
		G4	1540±23.81	3.74
4	College of Pharmacy	G5	860±15.34	2.09
		G6	1251.4±12.77	3.04
		P1	882.8±17.72	2.14
		P2	1244.2±58.21	3.02
		P3	830±22.68	2.01
		P4	384.2±17.26	0.93
5	The central Library	P5	1955.7±20.93	4.75
		P6	1178.5±12.88	2.86
		L1	2941.4±26.70	7.15
		L2	1647.1±19.96	4
		L3	1220±19.95	2.96
		L4	1012.8±11.51	2.46
6	College of Veterinary Medicine	L5	1447.1±24.00	3.51
		L6	1481.4±53.08	3.6
		M1	714.2±16.34	1.73
		M2	774.2±25.74	1.88
		M3	382.8±13.25	0.93
		M4	687.1±15.52	1.67

		M5	722.8±6.72	1.75
		M6	1257.1±23.05	3.05
7	College of physical education	R1	1897.1±31.40	4.61
		R2	487.1±15.58	1.18
		R3	1185.7±40.54	2.88
		R4	2157.1±59.60	5.24
		R5	2047.1±26.16	4.97
		R6	1564.2±11.17	3.8
8	College of Agriculture	F1	1630±48.95	3.96
		F2	1410±23.91	3.42
		F3	2051.4±16.57	4.98
		F4	2651.4±34.72	6.44
		F5	958.57±39.35	2.33
		F6	1580±28.31	3.84
9	Internal Sections for Girls	D1	2107.1±12.08	5.12
		D2	1257.1±12.98	3.05
		D3	828.5±41.02	2.01
		D4	1287.1±41.95	3.13
		D5	2270±32.51	5.52
		D6	1804.2±24.20	4.38
10	College of Administration Economics	E1	1825.7±25.74	4.43
		E2	1560±13.71	3.79
		E3	2548.5±19.73	6.19
		E4	881.4±26.01	2.14
		E5	320±9.73	0.77

		E6	631.4±25.90	1.53
11	College of Dentistry	C1	2128.5±11.79	5.17
		C2	1887.1±29.10	4.58
		C3	1645.7±42.84	4
		C4	1314.2±24.58	3.19
		C5	1778.5±22.95	4.32
		C6	707.1±25.97	1.71
12	College of Literature	V1	3007.1±12.13	7.31
		V2	1194.2±34.37	2.9
		V3	571.4±12.64	1.38
		V4	1064.2±23.76	2.58
		V5	1778.5±19.80	4.32
		V6	2175.7±13.87	5.29
13	College of Islamic Sciences	N1	1914.2±28.51	4.65
		N2	1598.5±18.76	3.88
		N3	2375.7±15.43	5.77
		N4	1828.5±30.05	4.44
		N5	2110±30.03	5.13
		N6	520±10.38	1.26
14	Printing	U1	1324.2±21.61	3.22
		U2	1821.4±16.31	4.42
Average			1431.25±646.2	3.48±1.57
The acceptable limit [8]				11.7ppm




Figure 6: Concentration of uranium in soil samples

4. Conclusions

- 1) There is a clear contrast between the uranium concentrations in samples, which is attributed to the fact that some soils are transferred soils, especially in college of (arts and sports), that we concluded that some soil are transfer from another places.
- 2) The current study found that the technique of counting the effects of fission fragments to calculate the concentration of uranium in soil samples using the nuclear track detector (CR-39) of the appropriate and good techniques to study the activity of radiation as it does not need an electronic system and electric energy.
- 3) The concentration of uranium in soil samples is found in sample (V1) and is equal to (7.31)ppm and this value is close to the permissible limit which is equal to (11.7 ppm) that we be leave future monitoring those level is necessary.
- 4) It is recommended to use different and new methods and periodic and seasonal tests because radiation pollution is

influenced by atmospheric factors and environmental variables in order to for monitor this places.

References

- [1] AL-Mahmood .A. AL-Karim.A " The Environment and Health Impact of Radioactive Materials" Proceedings of the Scientific Conference on the Impact of the Use of DU Weapons on Man and the Environment in Iraq, Part I, p. 15 (2002).
- [2] Dietz, L. A. "Uranium Battle Fields", Progressive Alliance for Community, New Mexico, College of Science City University Journal. (1993).
- [3] Fahad.A.A, Shehab.R.M, Abdel-Hussein Wens, Hossam El-Din Ahmed, Ali Abbas Mohamed, Mahmoud Shaker Mahmoud "Study of the movement and transfer of depleted uranium in the soils of the southern regions of Iraq" Proceedings of the Scientific Conference on the Impact of the Use of DU Weapons on Man and the Environment in Iraq, Part I, p. 180 (2002).

- [4] Al-Rahal, I. K. ((Calculation of Concentrations and Transfer Factor of Uranium in Phosphate Fertilizers from Soil to Plants using the Nuclear Impact Detector CR-39)). Master Thesis, Baghdad University, college of Science for Girls, (2008).
- [5] Saied1.B.M. Al-Khafaji2.R.M. Al-Bayati1.A.T. Measurement of Uranium Concentration in the Soil Samples by Using Solid State Nuclear Track Detectors (SSNTDs).Science and Technology . vol.2,(2016).
- [6] Rahim, Z. J. "Study of the Effect of High Pressure Lines on Radon Concentrations in the Air Using the Nuclear Impact Detector CR-39" Baghdad University, College of Education (Ibn al-Haytham), Master Thesis, (2009).
- [7] Tawfiq.N.F. H.Mansour.H.L. and Karim.M.S. "Measurement of Radon Gas Concentrations in Tap Water for Baghdad Governorate by Using Nuclear Track Detector (CR-39)", International Journal of Physics, Vol.3(6), (233-238). (2015).
- [8] United Nations Scientific Committee on the Effects of Atomic Radiation. UNSCEAR 2000 report to the general assembly, with scientific annexes.Sources and effects of ionizing radiation. United Nations, New York,(2000).

