

# Anthropometric Study of Knee Joint in Anterior Cruciate Ligament Insufficient Knees using Magnetic Resonance Imaging

Prathyush Chitithoti<sup>1</sup>, Vella Sandeep<sup>2</sup>, A. Srinivasa Rao<sup>3</sup>

<sup>1</sup>Junior Resident, Department of Orthopaedics, Katuri Medical College and Hospital, Guntur, India

<sup>2</sup>Assistant Professor, Department of Orthopaedics, Katuri Medical College and Hospital, Guntur, India

<sup>3</sup>Professor & Head of the Department, Department of Orthopaedics, Katuri Medical College and Hospital, Guntur, India

**Abstract:** ***Purpose:** The objective of this study was to evaluate morphological features on MRI in patients with ACL insufficient knees. **Methods:** Magnetic resonance imaging (MRI) of the knee of 40 anterior cruciate ligament sufficient patients and 20 anterior cruciate ligament intact patients who came to Katuri Medical College and Hospital, Guntur, between Sep 2017 and August 2018 were reviewed. Their intercondylar notch, femoral condyles dimensions and tibial slopes were measured using MRI. **Results:** The ACL insufficient knees and ACL intact knees were comparable in terms of intercondylar notch width. Lateral femoral condyle height and transepicondylar axis are more in ACL insufficient knees. Medial posterior tibial slope is more than Lateral posterior tibial slope in both ACL insufficient knees and ACL intact knees. **Conclusion:** The ACL insufficient patients had more Lateral femoral condyle width, intercondylar notch width, intercondylar notch height compared to ACL intact knees. ACL-ruptured patients had statistically significant smaller heights of femur's lateral condyle. MTS is more than LTS in both ACL insufficiency patients and ACL intact patients.*

## 1. Introduction

The dimensions of the intercondylar notch are such that there is little space not filled by the cruciate ligaments. It is this intimate relationship between the cruciate ligaments and the notch that may give rise to impingement of these structures following anterior cruciate ligament (ACL) reconstruction. Impingement of the ACL on the intercondylar notch will occur if the notch is stenotic, the ACL is relatively large, or a combination of both. This impingement will jeopardize the surgical repair.

Magnetic resonance imaging (MRI) has been proven as the gold standard method of choice for non-invasive evaluation of the ACL. Multiple studies have been executed in which measurements were taken on MR images to identify the characteristics of the ACL and its surroundings [1, 2]. Furthermore, MRI has been utilized during the evaluation of knee kinematics with and without an ACL rupture [3, 4].

To the best of the authors' knowledge, an assessment of multiple morphological features and their risk for ACL rupture has not been performed. Some studies have investigated the morphological features on the femoral side, while others have studied those on the tibial side. As such, the objective of this study was to evaluate multiple morphological features on both the femoral and tibial sides of the knee on MR imaging in patients with and without ACL rupture. In this way, it is possible to analyze the tibial and femoral morphological features both separately and combined and to evaluate a possible morphological relationship. The hypothesis of this study was that multiple individual and combined morphological features that confer a higher risk of sustaining an ACL rupture could be identified. This complete assessment of multiple morphological features and thereby the identification of

predisposing factors for an ACL rupture will identify the potential areas of interest for ACL injury prevention.

## 2. Materials and Methods

Magnetic resonance imaging scans from 20 patients with an intact ACL and 40 patients with a ruptured ACL who underwent an ACL reconstruction at Katuri Medical College and Hospital, Guntur, between May 2017 and April 2018 were prospectively studied. MRIs for non-injured knees were taken for knee complaints other than ACL injury. They were interpreted to have either meniscal injury or no identifiable pathology. Patients included in this study are of age 18 years to 60 years. Exclusion criteria for both groups included previous surgical interventions to the knee and patients with knee arthritis. The MR images used for this study were obtained with a 1.5 Tesla MRI with the knee in full extension. Slice thickness was 3 mm. Measurements were made in a standard, open-source image processing software for DICOM images.

In this study, the morphological features measured were the transepicondylar axis, lateral femoral condyle width, medial femoral condyle width, medial femoral condyle anteroposterior diameter, lateral femoral condyle anteroposterior diameter, the notch width, intercondylar notch height, medial tibial slope, the lateral tibial slope, medial tibial anteroposterior diameter, lateral tibial anteroposterior diameter.

### Transepicondylar axis:

The direction of axial slice imaging of the MRI is the slice perpendicular to the femoral mechanical axis in the coronal plane and perpendicular to the long axis of the femur in the sagittal plane. The axial slice on the most prominent part of both femoral condyles was selected for analysis. The transepicondylar axis (TEA) was defined as a line between

the most medial and the most lateral prominences of the epicondyles.



Figure 1: Transepicondylar axis

**Femoral condyle width, Notch width and Notch height:**

It is measured in coronal view of the knee joint MRI. A line is drawn at the most inferior part of the femoral condyles. A line parallel to this line at the level of the popliteal groove is shown, where the notch width (NW), medialcondyle width (MCW) and lateral condyle width (LCW) were measured. Femoral condyle anteroposterior diameter is measured at a section of axial slice where maximum anteroposterior diameter of lateral condyle is present. A horizontal line is drawn touching both the posterior surfaces of medial and lateral femoral condyles. Lateral femoral condyle Anteroposterior diameter (LAP) is measured from the anterior most part of the condyle to the previous horizontal line. Medial femoral condyle anteroposterior diameter (MAP) is measured from the anterior most part of the condyle to the previous horizontal line



Figure 2: Lateral femoral condyle width (LFCW), Medial femoral condyle width (MFCW)




Figure 3: Lateral femoral condyle Anteroposterior diameter (LAP), Medial femoral condyle Anteroposterior diameter (MAP)

Notch width is measured at three different locations in axial section where the popliteal groove appears maximum width. First measurement at the a line through the centre of the popliteal groove (NW - PG), second measurement at the level of two thirds of the notch height (NW -2/3), the third measurement at the level of the articular margins (NW - IAM) of the medial and lateral femoral condyles. The intercondylar notch height (NH) which was the perpendicular distance from the apex of the notch to the horizontal line at the level of the articular margins of the medial and lateral femoral condyles. These measurements are also noted two sections above and two sections below this level.




Figure 4: Intercondylar Notch Height (NH), Notch width at level of two thirds of the notch height (NW -2/3), Notch width -popliteal groove (NW - PG), Notch width at the level of the articular margins (NW - IAM)

**Lateral femoral condyle height:**

This is measured on sagittal sections of the MRI at the level of maximum diameter of lateral femoral condyle. Two circles were drawn, one anterior and one posterior, respectively. These matched the rounded anterior and posterior profile of the lateral condyle. A line is drawn at the maximum distance between the two circles touching the

anterior and posterior corticals. Two lines parallel to this line were drawn, proximal line touching the posterior circle, distal line touching the anterior circle. The distance between

these two lines is the measurement of height of the lateral femoral condyle (LFC-HT).




Figure 5: Height of the lateral femoral condyle (LFC-HT)

### Tibial plateau slope

Medial tibial plateau slope (MTPS), Lateral tibial plateau slope (LTPS) were measured using the method proposed by Hashemi et al.<sup>5</sup>




Figure 6: Lateral tibial plateau slope (LTPS), Medial tibial plateau slope (MTPS)

ACL insufficiency and ACL intact groups were compared using the 2-sample t test. A p value of <math><0.05</math> was considered statistically significant.

### 3. Results

The ACL-injured and ACL-intact groups were comparable in terms of the TEA (7.626 VS 7.4655,  $P = 0.363$ ), LCW (2.685 vs 2.277,  $p = <0.001$ ), MCW (2.3675vs2.3875,  $p = 0.21$ ), LAP (5.71575vs5.79,  $p = 0.011$ ), MAP


(5.4535vs5.616,  $p = <0.001$ ), At the level of maximum popliteal groove NW -2/3 (2.081 vs1.599,  $p = <0.001$ ), NW - PG (2.30125vs2.1525,  $p = <0.001$ ), NW - IAM (2.3745vs2.1165,  $p = <0.001$ ), NH (2.3955vs2.1075,  $p = <0.001$ ), 2 levels above NW -2/3 (1.655vs1.579,  $p = <0.001$ ), NW-PG (2.048vs2.033,  $p = 0.405$ ), NW - IAM (2.068vs2.0415,  $p = 0.161$ ), NH (2.166vs2.423,  $p = <0.001$ ), 2 levels below NW-2/3 (2.167vs1.896,  $p = <0.001$ ), NW -PG (2.207vs2.14,  $p = 0.255$ ), NW - IAM (2.344vs2.154,  $p = <0.001$ ), NH (2.12125vs1.822,  $p = <0.001$ ), LFC - HT

(3.497vs3.666, p= 0.07), LTPS (6.568vs2.025, p= <0.001), MTPS (9.208vs3.74, p= <0.001).

**4. Discussion**

Knee static stability is enabled by ligaments, menisci, joint capsule, and bony geometry, whereas knee dynamic stability is an integrated function of the neuromuscular system. Injury to the ACL can occur secondary to excessive valgus stress, forced external rotation of the femur on a fixed tibia, and hyperextension.<sup>6-8</sup> The most important finding of the present study is that there were bony morphological differences between ACL-insufficiency patients and ACL-intact subjects. Therefore, it is important to understand whether MRI measurements of bony morphology can be used reliably to determine a difference between ACL-injured patients and non-injured patients.

Transepicondylar axis is more in ACL Insufficiency patients 7.626cms when compared to ACL intact patients (7.465 cms). Lateral condyle width (LCW) is more in ACL Insufficiency patients (2.685 cms) when compared to ACL intact patients 2.277cms. Medial condyle width (MCW) is similar in both ACL Insufficiency patients 2.367 cms and ACL intact patients 2.387cms.




**Table 1:** TEA Width is more in ACL Injured knee in our study

Lateral femoral condyle anteroposterior diameter (LAP) is similar in both ACL insufficiency patients and ACL intact patients. Medial femoral condyle anteroposterior diameter (MAP) in ACL insufficiency patients (5.4535cms) is less than ACL intact patients (5.616cms).

At the level of maximum popliteal groove, Notch width at 2/3<sup>rd</sup> height (NW -2/3) is 2.081cms in ACL insufficiency knees which is more than ACL intact knees (1.739cms). Which is still more than Kelly klassen study 1.62cms. Notch width at the maximum popliteal groove level NW – PG is 2.261 cms is more in ACL Insufficient knees when compared to ACL intact knees (2.152 cms) which is even more than Kelly klassen study 2.14cms.

Notch width at the inferior articular margin level NW– IAM is 2.194cms is more in ACL Insufficient knees when compared to ACL intact knees (2.116cms) which is even more than Kelly klassen study 1.94cms.

Our study has measured the same parameters two sections above and two sections below the level of maximum popliteal groove. Two sections above the measures are similar to ACL insufficient knees to ACL intact knees. Two sections below, Notch width at 2/3<sup>rd</sup> height is more in ACL insufficiency patients and similar at popliteal groove and inferior articular margin.




**Table 2:** Notch Width (average of ACL insufficient knees and ACL Intact knees) is identical in both the studies.

Many studies have been performed previously to identify potential risk factors for ACL injury, and the femoral notch has been the main topic of interest. The majority of these studies have suggested that a smaller notch width and notch width index can be correlated to an ACL rupture<sup>(9-13)</sup>. Studies have also shown that women have smaller notch sizes than men. This may lead to the assumption that the smaller notch possibly contributes to a higher frequency of ACL ruptures in women<sup>(14-19)</sup>. Lombardo et al.<sup>(20)</sup> described that there is no differences between ACL-injured and non-injured basketball players with respect to Notch width. Our study states that notch width is more in ACL insufficiency patients than ACL intact patients.

Notch height at Maximum popliteal groove section is more in ACL insufficiency patients (2.395 cms) when compared to ACL intact patients (2.107 cms), two sections above ACL intact patients (2.423cms) have more values ACL insufficiency patients (2.166cms). Two sections below notch height is more in ACL insufficiency patients (2.333cms) when compared to ACL intact patients (1.822cms).

Lateral femoral condyle height is more in ACL intact patients (3.656cms) when compared to ACL insufficiency patients (3.497cms). In Margarida et al (2016) also LFC-HT is more in ACL intact patients (4.52 cms) when compared to ACL insufficiency patients (4.125 cms). It was found that ACL-ruptured patients had statistically significant smaller heights of femur’s lateral condyle.



**Table 3:** LFC Height is more ACL Intact knees-both studies

Lateral tibial plateau slope is more in ACL insufficient patients ( $6.568^0$ ) when compared to ACL intact patients ( $2.025^0$ ). Medial tibial plateau slope is more in ACL insufficient patients ( $9.208^0$ ) when compared to ACL intact patients ( $3.74^0$ ). In our study MTS is more than LTS both in ACL insufficiency patients and ACL intact patients. In Lazar et al (2008) study LTS is more in ACL insufficiency patients' and MTS is more in ACL intact patients.

The geometry of the proximal tibia, in particular the PTS, is considered a factor of ACL injury.<sup>(21-23)</sup> The PTS directly affects loading of the ACL during a compressive axial force and contributes to the propensity of the ACL to rupture. For every  $10^\circ$  increase in PTS, there is an associated 6-mm increase in tibial translation, and the Lachman test shows a 3-mm increase for every  $10^\circ$  increase in the tibial slope.<sup>20</sup> In a cadaveric study, a small increase in the tibial slope does not affect anteroposterior translation or in situ force in the cruciate ligaments. Nonetheless, increasing slope causes an anterior shift in the tibia resting position that is accentuated under axial loads. Altering the tibial slope may therefore be beneficial for ACL-deficient or posterior cruciate ligament-deficient knees to improve joint congruity.<sup>(24)</sup>

## 5. Conclusion

The ACL insufficient patients have more Lateral femoral condyle width, intercondylar notch width, intercondylar notch height compared to ACL intact knees. ACL-ruptured patients had statistically significant smaller heights of femur's lateral condyle. MTS is more than LTS in both ACL insufficiency patients and ACL intact patients.

## References

- [1] Frobell RB (2011) Change in cartilage thickness, posttraumatic bone marrow lesions, and joint fluid volumes after acute ACL disruption: a two-year prospective MRI study of sixty-one subjects. *J Bone Joint Surg Am* 93:1096–1103
- [2] Oldrini G, Teixeira PG, Chanson A, Erpelding ML, Osemont B, Louis M, Blum A (2012) MRI appearance of the distal insertion of the anterior cruciate ligament

- of the knee: an additional criterion for ligament ruptures. *Skeletal Radiol* 41:1111–1120
- [3] Espregueira-Mendes J, Pereira H, Sevivas N, Passos C, Vasconcelos JC, Monteiro A, Oliveira JM, Reis RL (2012) Assessment of rotatory laxity in anterior cruciate ligament-deficient knees using magnetic resonance imaging with Porto-knee testing device. *Knee Surg Sports TraumatolArthrosc* 20:671–678
- [4] Haugom BD, Souza R, Schairer WW, Li X, Ma CB (2012) Evaluating rotational kinematics of the knee in ACL-ruptured and healthy patients using 3.0 Tesla magnetic resonance imaging. *Knee Surg Sports TraumatolArthrosc* 20:663–670
- [5] Hashemi J, Chandrashekar N, Gill B, Beynon BD, Slauterbeck JR, Schutt RC Jr, et al. The geometry of the tibial plateau and its influence on the biomechanics of the tibiofemoral joint. *J Bone Joint Surg Am* 2008;90:2724–34.
- [6] Childs SG. Pathogenesis of anterior cruciate ligament injury. *OrthopNurs* 2002;21:35–40.
- [7] Hootman JM, Dick R, Agel J. Epidemiology of collegiate injuries for 15 sports: summary and recommendations for injury prevention initiatives. *J Athl Train* 2007;42:311–9.
- [8] Griffin LY, Agel J, Albohm MJ, Arendt EA, Dick RW, Garrett WE, et al. Noncontact anterior cruciate ligament injuries: risk factors and prevention strategies. *J Am AcadOrthoSurg* 2000;8:141–50.
- [9] Cha JH, Lee SH, Shin MJ, Choi BK, Bin SI (2008) Relationship between mucoid hypertrophy of the anterior cruciate ligament (ACL) and morphologic change of the intercondylar notch: MRI and arthroscopy correlation. *Skeletal Radiol* 37:821–826
- [10] Domzalski M, Grzelak P, Gabos P (2010) Risk factors for Anterior Cruciate Ligament injury in skeletally immature patients: analysis of intercondylar notch width using Magnetic Resonance Imaging. *IntOrthop* 34:703–707
- [11] LaPrade RF, Burnett QM 2nd (1994) Femoral intercondylar notch stenosis and correlation to anterior cruciate ligament injuries. A prospective study. *Am J Sports Med* 22:198–202
- [12] Shelbourne KD, Davis TJ, Klootwyk TE (1998) The relationship between intercondylar notch width of the femur and the incidence of anterior cruciate ligament tears. A prospective study. *Am J Sports Med* 26:402–408
- [13] Shelbourne KD, Facibene WA, Hunt JJ (1997) Radiographic and intraoperative intercondylar notch width measurements in men and women with unilateral and bilateral anterior cruciate ligament tears. *Knee Surg Sports TraumatolArthrosc* 5:229–233
- [14] Murshed KA, Cicekcibasi AE, Karabacakoglu A, Seker M, Ziylan T (2005) Distal femur morphometry: a gender and bilateral comparative study using magnetic resonance imaging. *SurgRadiolAnat* 27:108–112
- [15] Stijak L, Radonjic V, Nikolic V, Blagojevic Z, Aksic M, Filipovic B (2009) Correlation between the morphometric parameters of the anterior cruciate ligament and the intercondylar width: gender and age differences. *Knee Surg Sports TraumatolArthrosc* 17:812–817

- [16] Tillman MD, Smith KR, Bauer JA, Cauraugh JH, Falsetti AB, Pattishall JL (2002) Differences in three intercondylar notch geometry indices between males and females: a cadaver study. *Knee* 9:41–46
- [17] van Eck CF, Martins CA, Vyas SM, Celentano U, van Dijk CN, Fu FH (2010) Femoral intercondylar notch shape and dimensions in ACL-injured patients. *Knee Surg Sports TraumatolArthrosc* 18(9):1257–1262
- [18] Vrooijink SH, Wolters F, Van Eck CF, Fu FH (2011) Measurements of knee morphometrics using MRI and arthroscopy: a comparative study between ACL-injured and non-injured subjects. *Knee Surg Sports TraumatolArthrosc* 19:S12–S16
- [19] Wolters F, Vrooijink SH, Van Eck CF, Fu FH (2011) Does notch size predict ACL insertion site size? *Knee Surg Sports TraumatolArthrosc* 19:S17–S21
- [20] Lombardo S, Sethi PM, Starkey C (2005) Intercondylar notch stenosis is not a risk factor for anterior cruciate ligament tears in professional male basketball players: an 11-year prospective study. *Am J Sports Med* 33:29–34
- [21] Feucht MJ, Mauro CS, Brucker PU, Imhoff AB, Hinterwimmer S. The role of the tibial slope in sustaining and treating anterior cruciate ligament injuries. *Knee Surg Sports TraumatolArthrosc* 2013;21:134–45.
- [22] Hashemi J, Chandrashekar N, Mansouri H, Gill B, Slaughterbeck JR, Schutt RC Jr, et al. Shallow medial tibial plateau and steep medial and lateral tibial slopes: new risk factors for anterior cruciate ligament injuries. *Am J Sports Med* 2010;38:54– 62.
- [23] Khan MS, Seon JK, Song EK. Risk factors for anterior cruciate ligament injury: assessment of tibial plateau anatomic variables on conventional MRI using a new combined method. *IntOrthop* 2011;35:1251–6.
- [24] Giffin JR, Vogrin TM, Zantop T, Woo SL, Harner CD. Effects of increasing tibial slope on the biomechanics of the knee. *Am J Sports Med* 2004;32:376–82.