

# Effect of Metal Ions on Oil Degradation Activity of *Bacillus* and *Acinetobacter* Isolated from Oil Contaminated Sites of Mechanical Workshops

Dr. Praveen Reddy .P

Department of Microbiology, Vivekananda Degree and PG College, Karimnagar-505001, Telangana, India

**Abstract:** Used engine oils released from various automobile mechanical workshops pollute the environment. The various toxic chemical compounds present in the used engine oil cause deleterious effects to environment and living beings. In nature, there are some microbes which have the ability to degrade organic pollutants present in used engine oils. Such microbes can be isolated, identified and can be used for biodegradation of organic pollutants. In the present paper, effect of different metal ions on oil degradation activity of two bacterial species viz., *Bacillus* and *Acinetobacter* was determined using minimal salt broth supplemented with used engine oil and different metal ions. Majority of the metal ions selected for the study were found to be important for the enhancement of oil degradation activity of both the bacteria. Presence of only two ions viz.,  $\text{Fe}^{3+}$  and  $\text{Mn}^{2+}$  decreased the oil degradation ability of the bacteria.

**Keywords:** Used engine oil, organic pollutants, oil degradation, *Bacillus*, *Acinetobacter*, metal ions

## 1. Introduction

Automobile mechanical workshops are the main sources of used engine oils. Used engine oils contain heavy metals, wide variety of aliphatic and polyaromatic compounds. Used engine oils when disposed from mechanical workshops and other sources reach surroundings and cause environmental pollution. Especially, some of the polyaromatic compounds present in used engine oil are highly mutagenic and carcinogenic. Naphthalene, one of the most toxic polyaromatic compounds adversely affects the various human organs like kidney, heart, lungs etc. The chemical compounds present in the used engine oil change the physico-chemical conditions of soil and immobilize plant nutrients [1,2,3]. Some microorganisms in the environment have the ability to metabolize the hydrocarbons present in used engine oils. These microbes can be employed for the degradation of toxic hydrocarbons. The process of degradation of organic pollutants by employing microbes is called biodegradation. Microbes which can degrade organic pollutants include *Mycobacterium*, *Streptococcus*, *Bacillus* etc. Biodegradation process is the main technique employed in bioremediation [4]. In the present work twenty four oil degrading bacteria were isolated from oil contaminated soils of mechanical workshops of Autonagar, Kaman region of Karimnagar town, Telangana and identified till genus level. Thirteen were identified as *Bacillus* species, four were *Acinetobacter* species, three were *Pseudomonas* and remaining three were *Micrococcus* species. Bacterial species belonging to four genera (viz., *Bacillus*, *Acinetobacter*, *Pseudomonas* and *Micrococcus*) were isolated. In the present paper, two bacterial species each one from two bacterial genera (viz., *Bacillus* and *Acinetobacter*) exhibiting highest oil degrading activity were selected and effect of different metal ions on the oil degradation of activity these two bacteria was studied.

## 2. Materials and Methods

### Medium and metal ions used

Effect of different metal ions on oil degradation activity of *Bacillus* and *Acinetobacter* was carried out using minimal salt medium (MS medium) supplemented with used engine oil (5%) as sole carbon for the growth of bacteria [5]. Effect of seven metal ions on oil degradation activity of bacteria was determined. The metal ions selected were  $\text{Fe}^{3+}$ ,  $\text{Co}^{2+}$ ,  $\text{Cu}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Mn}^{2+}$ ,  $\text{MoO}_4^{2-}$  and  $\text{Zn}^{2+}$ . The MS medium with 5% used engine oil and all seven metal ions is considered as negative control and MS medium with 5% used engine oil without any ions as positive control. The effect of absence of each ion on oil degradation ability of bacteria was studied in the presence of remaining all ions in the medium [6].

### Inoculum preparation

The bacterial cultures were inoculated separately into 10 ml of MS broths supplemented with used engine oil (5%) as sole carbon source and incubated at  $30^\circ\text{C}$  for one day. After one day each bacterial culture broth is subjected to centrifugation at 3500 rpm and cell pellets of bacteria were obtained. The bacterial cell pellets were washed and grown in MS broth supplemented with used engine oil till the optical density (OD) of each culture was read 1.0 at 600nm in colorimeter [7.] Each such 0.5 ml of bacterial culture was used as inoculum and transferred to 10 ml of MS medium (with used engine oil and different metal ions combinations prepared separately) and incubated for seven days. After incubation period the OD of the bacterial cultures biomass was measured at 600 nm.

The growth of bacteria was measured in terms of OD of their cell mass in culture broth. As the broth contains used engine as the only source of carbon the growth of bacteria is the direct measure of oil degradation activity. All the experiments were carried out with triplicates and standard deviation (SD) value was calculated for each result.

### 3. Results and Discussion

In the present work effect of absence of selected metal ions on biodegradation ability (in terms of growth) of *Bacillus* and *Acinetobacter* was carried out. Majority of the cellular enzymes activity is under the influence of metal ions. Metal ions act as enzyme activators. Metal ions change flow of electrons in either substrates or enzymes and control the biocatalysis [8]. On the other hand some metal ions inhibit the activity of certain enzymes. One of the key roles of certain metal ions is involvement in the formation of protein synthetic machinery [9]. In the absence of  $\text{Fe}^{3+}$  and  $\text{Mn}^{2+}$  ions the biodegradation ability (growth) of the bacteria increased when compared to growth observed in negative control and positive control (Table-1). Jernejc and Legisa (2002) reported that  $\text{Fe}^{3+}$  inhibited the activity of malic dehydrogenase [10]. In *Lysobacter enzymogenes*  $\text{Mn}^{2+}$  ions (at concentration more than 0.01 mM) inhibited the formation of nuclease, RNAase and alkaline phosphatase [11]. In the absence of  $\text{Ca}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Cu}^{2+}$  and  $\text{MoO}_4^{2-}$  ions the bacteria showed decreased oil degradation activity (growth) (Table-1). In *Myriococcus thermophilus* and *Humicola insolens* the  $\text{Ca}^{2+}$  ions enhanced the cellobiose dehydrogenase activity [12]. In the presence of  $\text{Co}^{2+}$  ions the activity of alkaline phosphatase was increased in *Bacillus licheniformis* [13].  $\text{Cu}^{2+}$  ions (in the form of  $\text{CuSO}_4$ ) increased the activity of ascorbate oxidase in callus of Cucumber [14]. Al-Issawi reported from his studies on two wheat cultivars that presence of  $\text{MoO}_4^{2-}$  ions had enhanced the activity of antioxidant enzymes [15]. In both the bacteria the growth was adversely affected in the absence of  $\text{Zn}^{2+}$  ions in the medium (Table-1). Similar results were obtained for Alves (2007) while working on dibenzothiophene desulfurizing *Gordonia alkanivorans* strain 1B [6].

**Table 1:** Effect of Metal ions on growth (oil degradation activity) of *Bacillus* and *Acinetobacter*

Sl. No	Metal ion	OD (Growth ) of <i>Bacillus</i> at 600 nm	OD (Growth ) of <i>Acinetobacter</i> at 600 nm
1	Negative control (contains all ions)	3.35 ± 0.14	3.03 ± 0.05
2	Positive control (contains no ions)	1.81 ± 0.09	1.60 ± 0.12
3	$\text{Fe}^{3+}$ absence	3.75 ± 0.08	2.40 ± 0.08
4	$\text{Co}^{2+}$ absence	3.05 ± 0.05	2.73 ± 0.05
5	$\text{Cu}^{2+}$ absence	2.61 ± 0.12	3.17 ± 0.14
6	$\text{Mn}^{2+}$ absence	3.65 ± 0.17	2.60 ± 0.09
7	$\text{MoO}_4^{2-}$ absence	2.91 ± 0.09	3.27 ± 0.05
8	$\text{Zn}^{2+}$ absence	3.35 ± 0.14	1.83 ± 0.17
9	$\text{Ca}^{2+}$ absence	1.81 ± 0.05	2.93 ± 0.05

### 4. Conclusion

The oil degrading bacteria *Bacillus* and *Acinetobacter* exhibited decreased oil degradation activity (growth) in the absence of  $\text{Ca}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Cu}^{2+}$ ,  $\text{MoO}_4^{2-}$  and  $\text{Zn}^{2+}$  ions indicating their role in growth and metabolic activities. Only two metal ions viz.,  $\text{Fe}^{3+}$  and  $\text{Mn}^{2+}$  decreased the growth of both the oil degrading bacteria. Overall,  $\text{Zn}^{2+}$  ions can be regarded as key metal ions for growth and various cellular activities as its absence adversely affected the oil degradation activity (growth) of both the bacteria.

### References

- [1] Ugoh SC and Moneke L, 2011. Isolation of bacteria from Engine Oil Contaminated Soils in Auto-mechanic Workshops in Gwagwala da Abhuja, FCT-Nigeria. Academia Arena 3(5), 28-33.
- [2] Mbachu AE., Onochie CC., Agu KC., Okafar OI., Awah NS, 2014. Hydrocarbon degrading potentials of indigenous bacteria isolate from Auto-mechanic Workshops at MGBUKA-NKPOR, Nigeria. Journal of Global Biosciences. 3(1), 321-326.
- [3] Jahir AK and Syed HAR, 2011. Isolation and Characterization of micro-organisms from oil contaminated sites. Advances in Applied Science Research. 2(3), 455-460.
- [4] Ogunbayo AO., Bello RA., Nwagbara U, 2012. Bioremediation of engine oil contaminated site. Journal of Emerging Trends in Engineering and Applied Sciences. 3(3), 483-489.
- [5] Nwinyi OC., Ajaja O., Nwinyi, 2014. Growth dynamics of bacteria isolated from spent engine oil contaminated tropical soil. Research Journal of Environmental and Earth Sciences. 6(9), 430-436.
- [6] Alves LMG, 2007. Dibenzothiophene desulfurization by *Gordonia alkanivorans* strain 1B. University of Libson, Libson, Portugal (Ph.D Thesis).
- [7] Thenmozhi R., Nagasathya A., Thajuddin N, 2011. Studies on biodegradation of used engine oil by consortium cultures. Advances in Environmental Biology. 5(6), 1051-1057.
- [8] Glusker JP., Katz AK., Block CW, 1999. Metal ions in Biological Systems. The Rigaku Journal. 16(2), 8-16.
- [9] Joshi PA., Singh S., Shekhawat DB, 2015. Effect of metal ions on growth and biosurfactant production by Halophilic bacteria. Advances in Applied Science Research. 6(4), 152-156.
- [10] Jernejc K and Legisa M, 2002. The influence of metal ions on malic enzyme activity and lipid synthesis in *Aspergillus niger*. FEMS Microbiology Letter. 217 (2), 185-190.
- [11] Tigerstrom RGV, 1983. The effect of magnesium and manganese ion concentrations and medium composition on the production of extracellular enzymes by *Lysobacter enzymogenes*. Microbiology. 129, 2293-2299.
- [12] Sculz C., Ludwig R., Micheelsen PO., Silow M., Toscano MD., Gorton L, 2012. Enhancement of enzymatic activity and catalytic current of cellobiose dehydrogenase by calcium ions. Electrochemistry Communications. 17, 71-74.
- [13] Spencer DB., Chen CP., Hulett FM, 1981. Effect of cobalt on synthesis and activation of *Bacillus licheniformis* alkaline phosphatase. Journal of Bacteriology. 145(2), 926-923.
- [14] Sekiya J., Mizuno K., Kimura O., Shimose N, 1990. Ascorbate oxidase in Cucumber Calli and enhancement of enzyme activity by copper sulfate. Soil Science and Plant Nutrition. 36(1), 43-51.
- [15] Al-Issawi M., Rihan HZ., Al-Shmgani H., Fuller MP, 2016. Molybdenum application enhances antioxidant enzyme activity and COR15a protein expression under cold stress in wheat. Journal of Plant Interactions. 11(10), 5-10.