

The Rational Use of Antibiotics

Sadhika

Abstract: ***Introduction:** The abuse of antibiotics as well as the development of microbial resistance in them is a global phenomenon. **Aim:** The purpose of this study is to highlight the rational use of antibiotics and limitation of microbial resistance. **Materials and methods:** Extensive review of the recent literature was conducted in database Medline and via the link of the (HEAL-Link) Greek Association of Academic Libraries, using key words: antibiotics, medicine and rational use of antibiotics. **Results:** The rational use of antibiotics that should be random. It should be thought and should be based on guidelines. The accurate diagnosis, the patient's condition, the site of infection, the severity of microbial cause and its sensitivity to antibiotics, the pharmacokinetics and pharmacodynamics of antimicrobials, their side effects and costs are the main elements which must be supported in every aspect for the use. **Conclusion:** Physician and other health care professionals should prescribe antibiotics only when necessary and based on condition severity, according to the existing guidelines.*

Keywords: Antibiotics, Medicine and Rational use of antibiotics

1. Introduction

The overuse of antibiotics as well as the development of resistance in microbes is a global phenomenon. It concerns both the World Health Organization (WHO) and individual countries [1]. There are documented studies that support, beyond doubt, that restricting the misuse of antibiotics reduces resistance [2]. They all agree though, that the only way to reduce the resistance of antibiotics is through proper use and reducing abuse. We also all agree that in order to reduce the overuse of antibiotics simultaneous training is needed for both doctors and patients. Studies showed that unilateral intervention either by patients themselves or physicians themselves, have substantial results. So, it is now clear that the only way for the proper use of antibiotics is through intervention programs by both doctors and patients [3]. It is concluded that only by proper educational intervention for both doctors and patients by the right people with the right tools, can improve the problem of abuse of antibiotics sufficiently and to eliminate the risk of antimicrobial resistance gradually which insidiously results in major medical and social problem. Yet, times have changed and we now need to retrain patients and physicians about the proper use of antibiotics for the health of their children and their own [4]. The purpose of this study is to highlight the rational use of antibiotics to achieve optimal outcome of the infection with minimization of toxicity, reduction of costs and limitation of microbial resistance.

2. Material and Methods

Extensive review of the recent literature was conducted in electronic database Medline and via the link of the Greek Association of Academic Libraries (HEAL-Link), using the following key words: antibiotics, medicine, misuse and impact.

Antibiotics

Antibiotics are substances that destroy the virus without harming the host, human. Etymologically, the word comes from the anti-biotic and means anti-against and biotic-used for life [5]. Antibiotics are either natural substances that are produced in nature by microorganisms or synthetic substances, which have been prepared in the laboratory. To be considered a clinically effective antibiotic and therefore useful in medicine, the destruction or growth inhibition of

the microorganism is achieved in the respective concentrations of the antibiotic in the body [6]. To date, at least 4,000 antibiotics have been isolated from cultures of microbes and 30,000 have been prepared semisynthetic. In every day practice, however, only 100 of these are used. Apart from Medicine, they are used in Agriculture, Veterinary and Food Chemistry [7].

Antibiotics may be [8]:

- Wide spectrum kill many types of bacteria e.g. penicillin. The broad spectrum antibiotics are active against many types of microbes such as bacteria, rickettsia, mycoplasmas, protozoa, and spirochetes.
- Narrow spectrum which kill certain types of bacteria e.g. isoniazid and should be used where possible to reduce the risk of colonization and super infection with resistant bacteria. Basically, antibiotics are classified as bactericides that kill germs, or bacteriostatic, preventing the growth of bacteria. These classifications are based on laboratory behavior of antibiotics, but in fact, both of these groups are able to treat a bacterial infection [9].

There are a number of factors which should be considered for each host when administering antibiotic formulations [10].

Genetic factors such as lack of G6PD:

- Renal and hepatic impairment are factors that will determine the type of antibiotic.
- Bactericidal drugs are necessary for life threatening infections such as sepsis.
- Diseases such as tumours, lymphomas, vasculitides and systemic diseases.
- Infections where the administration of antibiotics should be immediate because the presence of infection is life threatening. Such examples include meningitis, bacterial endocarditis, leukopenia and acute necrotizing cellulitis.

Before the administration of antibiotics, anaerobes aerobic cultures should be taken which must be monitored because after the initial treatment, failure will therefore change initial therapeutic regimen [11].

The Use of Antibiotics

From 2008, at European level, the 18th November was established as Awareness Day for Rational Use of

Antibiotics, and efforts are made using campaigns aimed at rationalizing and reducing the over-consumption of antimicrobial agents. Recipients of these messages are the general public to raise awareness and to reduce the consumption of antimicrobials, but above all, for health professionals at all levels of care [22]. The rational use of antibiotics is therefore, like any other therapeutic intervention in daily practice and it should not be random. It requires reflection and thought and should be based on rules. The correct diagnosis, the patient's condition, the location of the infection, the severity of the microbial cause sensitivities to antibiotics, the pharmacokinetics and pharmacodynamics of antimicrobials, the side effects and cost are the main elements which must be supported in every decision for their use [3].

Health professionals also need to bear in mind the following basic rules for the rational management of antibiotics [27]:

- a) Too many antibiotics together may act synergistically rather than competitively. The majority of infections can be treated with an antibiotic. But there are cases where it is necessary to combine the administration of antibiotics. This is the case where a polymicrobial infection. For examples intraventricular brain abscesses and infections of the lower limbs of diabetic patients [28].
- b) The use of many antibiotics results in more side effects. Adverse reactions to antibiotics are common and are divided into idiosyncratic, allergic and dose-dependent. Examples of serious allergic reactions are immediate hypersensitivity reaction to penicillin (laryngeal edema, circulatory collapse). When a patient taking many antibiotics had an allergic reaction, then all antibiotics are suspect and cannot reuse [3].
- c) The use of many antibiotics increases the risk of infection with resistant micro-organisms. When many antibiotics are administered to a patient, the spectrum of microorganisms killed is increased. The destruction of the flora of the oropharynx and gastrointestinal tract are harmful to humans. The micro-organisms that are normally found in these regions are competing to develop more resistant micro-organisms. The elimination of the normal flora of the indiscriminate use of antibiotics has resulted in the development of multiresistant pathogenic microorganisms [29].

Also, you shouldn't [30]:

- Allow people take antibiotics on their own initiative or at the instigation of others or primarily, if not prescribed by your doctor.
- The prescription of antibiotics is not necessary in every condition. If the doctor does not prescribe antibiotics do not deny that you're sick.
- The antibiotics to viral infection do not reduce the duration of symptoms or protect people around you to be none afflicted.
- When assumed by your doctor that it is necessary to administer antibiotics, they should be strictly adhered to dosage instructions and duration of your treatment.
- Follow the physician's instructions carefully in the selection of antibiotics or give the same medicine to your loved ones on your own [31].

The strategy of administration of the antibiotics [23, 24]:

- Patients with serious bacterial infections, the number of white cells are increased and even exceed granulocytes.
- Based on the clinical picture, laboratory tests can identify the anatomical location of the infection. For example, the combination of fever, burning on urination, and increased in number of pyosfairion urinalysis indicates infection [6].
- Patients who develop an infection during their hospital stay empiric antimicrobial therapy administration should take account of the sensitivity of the flora, not only the hospital and the department (e.g., ICU, surgery, clinical pathology) which developed the infection [7].
- The remarkable ability of the micro-organisms to adapt to any environment is that each new pathogenic is resistant to antibiotics administered in the previous period.
- Patients with meningitis should not be administered antibiotics which do not pass the blood brain barrier, such as the first generation cephalosporin, gentamicin, and clindamycin. In endocarditic the tag is protected from the action of antibiotics. For this reason, the treatment should be done with bactericidal drugs at high doses and for prolonged periods [3].
- The foreign body infections, prosthetic valves, prosthetic joints, pacemakers, it is almost impossible to eradicate without removing the foreign body.
- In acute infection, patients with neutropenia, have an increased mortality from sepsis and appropriate direct intravenous therapy (taking the appropriate cultures) with broad spectrum bactericidal antibiotics at high doses.
- Older people metabolize and excrete antibiotics at a slower pace. Therefore, there should be longer intervals between the doses of antibiotics. At this age, drugs with greater toxicity, such as aminoglycosides (nephrotoxicity and ototoxicity) should be avoided [25,26].

3. Conclusion

Antibiotics are an essential tool of medical use in common medical procedures, such as transplantation and chemotherapy. However, over the years, bacteria have acquired resistance to antibiotics. Resistant bacteria can be transmitted from animals to humans through the food chain or by direct contact. Many bacterial infections are becoming resistant to the treatments most commonly prescribed antibiotics [32]. The resistance of pathogenic microorganisms to antibiotics not only a problem for the patient, but also for the environment as it is populated by the same pathogen and are more likely to be affected by it. So doctors and other professionals should prescribe antibiotics only when necessary, based on existing guidelines.

References

- [1] World Health Organization (2000) WHO Global principles for the containment of antimicrobial resistance in animals intended for food: Report of a WHO consultation. Geneva.

- [2] Kyriopoulos J (2014) Introduction to the politics and economy of the medicine. Presentation of NSPC, National School of Public Health, Europe.
- [3] Kotsiftopoulos C, Kourkouta L, Papageorgiou M (2014) The use of antibiotic medicine. Monograph, Lap Lambert Academic Publishing, Saarbrucken, Germany.
- [4] Hellenic Centre for Diseases Control and Prevention (HCDCP) (2007) Guidelines for diagnosis and experiential therapy infections, Athens, Greece.
- [5] <http://www.lexigram.gr/lex/omor/>
- [6] Iosifidi H (2013) Relationship of consumption of antimicrobial medicinal products and microbial resistance in a third-hospital hospital. Study of the effect of therapeutic intervention instructions. Aristotle University of Thessaloniki, Thessaloniki, Greece.
- [7] Partalidou DA (2016) The global phenomenon of antimicrobial resistance. Presentation of a phenomenon. Thesis. Intermediary Program of Postgraduate Studies. Legal and Theological Schools. Department of Medicine-Dentistry. Aristotle University of Thessaloniki, Thessaloniki, Greece
- [8] Marcelos M (2006) Medical Pharmacology Teaching Notes. University of Ioannina, Ioannina, Greece.
- [9] Kousoulakos X, Fragoulakis B (2007) The Drug Market in Greece, Annual Report 2007. Health Insurance Observatory, Athens, Greece.
- [10] http://school.med.uoa.gr/Documents/3etos/Farmakologi_a%20II.pdf
- [11] EOF (2014) Primary care infections and their treatment. Athens.
- [12] Livermore DM, Brown DF (2001) Detection of beta-lactamase-mediated resistance. *J Antimicrob Chemother* 48: 59-64.
- [13] <http://www.who.int/mediacentre/factsheets/fs194/en/>
- [14] Ministry of Health and Social Solidarity (2008) National Action Plan to address of microbial endurance in antibiotics and of Infections in spaces provision of Health Services 2008-2012. Athens, Greece.
- [15] The World Health Organization (WHO) (2014) Antimicrobial resistance. Fact sheet N°194".
- [16] Batopoulos A (2007) Antimicrobial resistance to antibiotics. "An important unknown public problem health. National School of Public Health.
- [17] Cosgrove SE (2006) The relationship between antimicrobial resistance and patient outcomes: mortality, length of hospital stay, and health care costs. *Clin Infect Dis* 42: S82-S90.
- [18] Koukos G (2014) Microbial resistance to health and inflammation of periodontal disease and transfusional tissues. Aristotle University of Thessaloniki, Thessaloniki, Greece.
- [19] Strateva T, Yordanov D (2009) *Pseudomonas aeruginosa*-a phenomenon of bacterial resistance. *J Med Microbiol* 58: 1133-1148.
- [20] Ministry of Health and Social Solidarity (2008) End-of-file (EOF). For the proper use of these EOF. Antibiotics, EOF Publications, Athens, Greece.
- [21] Pratti A, Karanika M, Maniatis AN, Petinaki E, Spiliopoulou I, et al. (2007) Activity of linezolid against Gram-positive cocci: a multicentre study in Greek hospitals. *Int J Antimicrob Agents* 29: 604-605.
- [22] <http://www.onmed.gr/ygeia-eidhseis/story/349561/keelpno>
- [23] Giamarelou E (1988) Principles of antimicrobial chemotherapy of infections. *Med J* 53: 631-636.
- [24] Pafitou EN (2012) National Strategy of Cyprus for antibiotic resistance. Democracy of Cyprus. Ministry of Health. Nicosia, Greece.
- [25] Kourkouta L (2016) PolyPharmacy in Elderly. *JPPS* 103: 1-5.
- [26] Kourkouta L, Frantzana E, Iliadis Ch, Monios A (2016) Health Problems of the Elderly. Monograph. Scholar's Press. Saarbrucken, Germany.
- [27] Chrysos G (2017) Development of effective hospital-based Antibiotic Stewardship Program. The role of Infectious Disease Specialist. *Sci Chronicles* 22: 44-50.
- [28] Droutsa K (2008) Multifarmacy. Athens, Greece.
- [29] Abriouel H, Omar NB, Molinos AC, López RL, Grande MJ, et al. (2008) Comparative analysis of genetic diversity and incidence of virulence factors and antibiotic resistance among enterococcal populations from raw fruit and vegetable foods, water and soil, and clinical samples. *Int J Food Microbiol* 123: 38-49.
- [30] Gitonas M, Karabli EL (2006) Market Regulation Policies Drug. Papazisis, Athens, Greece.
- [31] Gioton M, Kyriopoulos G (2006) Politics and economy of the drug in Greece. Papazisis, Athens, Greece.
- [32] Spyraakis X (2006) Principles of public health in drug consumption. Athens, Greece.