

# An Analysis of Specific Acoustic Impedance, Apparent Molal Volume and Transport Properties of Non-Aqueous Solutions of Peptides

R.Padmavathy<sup>1</sup>, K.Dhanalakshmi<sup>2</sup>, E.Jasmine Vasantha Rani<sup>3</sup>, S.Sujatha<sup>4</sup>

<sup>1,3</sup>Associate Professor, Department of Physics, Seethalakshmi Ramaswami College,

<sup>2</sup>Research Scholar, Department of Physics, Seethalakshmi Ramaswami College,

<sup>4</sup>Assistant Professor, Department of Physics, Srimati Indira Ganthi college, Tiruchirappalli-620002, Tamilnadu, India

**Abstract:** Ultrasonic investigations in non-aqueous solutions of peptides provide useful information in understanding the Physico-chemical properties of the interacting components. Peptides and amino acids are used as probe molecule to understand the complex nature of protein. It is recognized that the study of the behavior of model compounds of proteins like amino acids and peptides as solutions help in understanding the factors governing the thermodynamic stability of the native structure of proteins. The analysis of ultrasonic and thermodynamic properties of proteins in non-aqueous solvent has been the area of interest. In the present investigation ultrasonic velocity, transport properties and other related parameters were successfully employed at different temperatures and concentrations to understand the structural changes occurring in the solutions.

**Keywords:** Ultrasonic velocity, transport properties, molecular interactions.

## 1. Introduction

Ultrasonic studies are extensively used to estimate the thermodynamic properties and to predict the intermolecular interactions in liquids [1]. It is most preferred in many fields such as pharmaceutical industry, biomedical research, chemical industry, water research and scattering spectroscopy, etc., [2]. Proteins are complex molecules, their behavior in solutions are governed by a combination of many specific interactions; it is a good approach to study simpler model compounds such as amino acids and peptides which are basic building blocks of the proteins [3]. Internal pressure and free volume which are the fundamental properties of the liquid state has been studied initially by Hildebrand et al [4-5] and subsequently by several workers [6-11]. The study of transport properties of liquid system has gained much importance during recent years [12] the molecular interaction in liquid solution can be studied by transport properties. The ultrasonic velocity and its derived parameters provide fruitful information regarding the nature of intermolecular interactions. In the present work an attempt has been made to investigate the behavior of peptide solution to study the molecular interaction through specific acoustic impedance, apparent molal volume and internal pressure and free volume at different temperatures (5°C to 55°C) & at various concentrations.

## 2. Material and methods

Ultrasonic investigations of peptide salt in formamide are carried out for five different concentrations from a low temperature to a high temperature. Basic parameters such as density, viscosity, ultrasonic velocity are determined with the help of constant temperature bath. Densities of the

solutions are measured through 25ml specific gravity bottle. Cannon Fenske viscometer is used to measure the time flow of the solution in order to calculate viscosity. Ultrasonic velocity is measured through ultrasonic interferometer of frequency 2MHz. the derived parameters such as transport properties ( $\pi_i$ ,  $v_f$ ) specific acoustic impedance ( $z$ ), apparent molal volume ( $\phi_v$ ) are determined using the basic parameters. The formulae used to calculate them are given below

$$\pi_i = bRT [K\eta/u]^{1/2} [\rho^{2/3}/M_{\text{eff}}^{7/6}] \quad \text{-----(1)}$$

$$V_f = [M_{\text{eff}} U/K\eta]^{3/2} c c \quad \text{-----(2)}$$

$$z = \rho u \quad \text{-----(3)}$$

$$\phi_v = 1000/C_1 \rho^0 (\rho^0 - \rho) + M/\rho^0 \text{ (ml/mol)} \quad \text{-----(4)}$$

## 3. Results And Discussion

### 3.1 Transport properties

Cohesive energy is the measure of internal pressure. In peptide salt, the internal pressure exhibits an increasing trend with respect to increasing molalities as shown in table and figure (1). The increase in cohesive energy indicates that there is a strong interaction prevails between the solute and solvent. Hence, this increase may be attributed to the structure making nature of the solute in the solvent [13-15].

Free volume is the free space available for the molecules to move [16]. The structure making and breaking effects of the samples are also confirmed by decreasing and increasing values of free volume with respect to molalities and temperatures as expected [17,18]. The variations of free volume with respect to molalities and temperatures are also confirm that there is strong solute-solvent interaction in peptide solutions [19].




Figure 1: Internal Pressure (atms)

Table 1: Internal Pressure (atms)

Molality	5°C	15°C	25°C	35°C	45°C	55°C
.001	19130	16330	13669	13147	11562	10586
.005	19393	16371	13736	13174	11720	10625
.01	19642	16389	13905	13254	12101	10787
.015	19647	16426	14044	13296	12029	10886
.02	19929	16911	14391	13691	12151	11085

.01	18.73	18.35	18.08	17.80	17.44	17.12
.015	18.77	18.40	18.12	17.82	17.50	17.14
.02	18.82	18.44	18.15	17.88	17.54	17.17




Figure 2: Free Volume (cc)

Table 2: Free Volume (cc)

Molality	5°C	15°C	25°C	35°C	45°C	55°C
.001	0.0087	0.0154	0.0288	0.0353	0.0562	0.0791
.005	0.0083	0.0153	0.0284	0.0351	0.0539	0.0782
.01	0.008	0.0152	0.0274	0.0344	0.0489	0.0747
.015	0.0081	0.0151	0.0265	0.0341	0.0497	0.0726
.02	0.0077	0.0139	0.0246	0.0311	0.0482	0.0687




Figure 4: Apparent Molal Volume (ml/mol)

Table 4: Apparent Molal Volume (ml/mol)

Molality	5°C	15°C	25°C	35°C	45°C	55°C
.001	1965.67	2231.57	596.97	-340.40	861.10	2306.06
.005	353.11	318.75	19.28	-133	221.68	453.17
.01	153.47	233.70	65.95	26.46	199.86	302.53
.015	125.12	201.97	88.39	66.48	182.23	257.14
.02	141.21	144.44	114.38	91.32	160.14	225.34




Figure 3: Specific Acoustic Impedance (RAYL)

Table 3: Specific Acoustic Impedance (RAYL)

Molality	5°C	15°C	25°C	35°C	45°C	55°C
.001	18.64	18.30	17.99	17.74	17.36	17.06
.005	18.68	18.33	18.05	17.79	17.41	17.10

### 3.2 Specific Acoustic Impedance

The variations of specific acoustic impedance with respect to the molality are given in table and in figure (1.3). The linear variations of specific acoustic impedance with concentrations and temperature may be due to the effect of solute-solvent interactions [19].

### 3.3 Apparent Molal Volume

The decrease in apparent molal volume with an increase in molality is observed in both the solutions which suggest that the strong ion-ion interaction [20] taking place in the solutions. The higher values of  $\phi_{\text{v}}$  suggest that there is strong solute-solvent interaction and lower values support the strong ion-ion interactions [21,22].

## 4. Conclusion

Ultrasonic technique is very useful in finding out the existence of interactions between the components of solutions. Dipeptides have been used in study of protein

structure and functions. Peptides and proteins are the most characteristic chemical compounds found in the living cells. Phenylalanine is an essential amino acid, it contains an aromatic side chain. Which contribute to ultraviolet absorbance and fluorescence properties. It is used in the manufacture of food & drink products and sold as nutritional supplement for its reputed analgesic & antidepressant effects. In the present work  $\pi_i, v_f, Z, \phi_v$  study have been carried out. The behavior of the solute in the solvent reveals that the salt act as structure maker.

## References

- [1] Palaniappan L and Karthikeyan V., Indian J. phys., 2005, 79 (2), 155
- [2] Ahmed Eid Fazary, Aly Fahmy Mohammed and Watalyia sb. Lebedeva., J. Chem. Thermodyn., 38 (2006) 1467-1473
- [3] J.H. Hildebrand and R.L.Scalt, Regular solution (prentice – Hall, New Jersey , 1962)
- [4] J.H. Hildebrand and R.L.Scalt, ed.3 Solubility of non – electrolytes (Reinhold, New York, 1964)
- [5] A.F.M. Borton, J. Chem. Educ. 48, 156 (1971)
- [6] C.V. Suryanarayana, Indian J. Pure Appl. Phys. 27, 751 ( 1989)
- [7] Digby D Macdonald and J.B. Hyne, can. J. Chem. 49-611(1971)
- [8] M.R.J. Dack, Aust. J. chem. 28,1643 (1975)
- [9] M.R.J. Dack, J. chem. Soc. Rev. (GB) 4,211(1975)
- [10] S.V. Subramanyam, T. Ramanujappa and E.S. Raja Gopal, Acustica 52,125 (1983)
- [11] E. Jr Acree William, Thermodynamic properties of non-electrolyte solutions (Academic press, Orlando, Florida., 1984) ch.8
- [12] Saraniya M.R and Prasaniya P.H., J.Pure and Applied Ultrasonics, 22 (2000) 111-114.
- [13] Suryanarayana C.V, Ind. J. Pure and Applied Physics, Vol.27, (1989) 751-757.
- [14] Beulah mary A, Jesintha Rakini J.H, Bharathi G., J. Acous. Soc. Ind., Vol 36 (2002) 14-19.
- [15] Bhatti S.S., J. Pure and Applied Ultrasonics., Vol 3, No.3 (1981) 45
- [16] Godhani D.R., etal., J. Pure and Applied Ultrasonics, Vol 23 (2001) 58-62.
- [17] Thabane V.A, Sangeetha Agarwal and Revetkar K.G., J. Acous. Soc. Ind., Vol 28, No.4 (2000) 305-308.
- [18] Sujatha.S, Padmavathy. R, Jasmine Vasantha Rani.E, International Journal of Pure and Applied physics 3(3)(2013)16-19.
- [19] Syal V.K.et.al., J.Acous.Soc. Ind., 30 (2002).
- [20] Sathyavathi A.v.Acourstica,28 (1973) 177.
- [21] Balachandran, Nature, 191 (1973) 164.
- [22] Dhanalakshmi, Rani G. and Arivadainambi T.J. Acous.Soc Ind Vol23, 1-4 (2000) 385-388.