Studies of Acoustic and Thermodynamic Properties of Electrolytes in Double Distilled Water at 303.15K

S. Mirikar, P. Pawar, G. Bichele

Department of Physics, Dayanand Science College, Latur, 413512(MS), India.

Departments of Physics, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad (M.S), India.

Abstract: The ultrasonic velocity, density and viscosity at 303.15K have been measured. for electrolytes (NaCl and Mgcl₂) in double distilled water at 303.15K. From experimental data acoustical parameters such as adiabatic compressibility (β_a), acoustic impedance (z), intermolecular free length (L_f), relative association (R_A) have been estimated using the standard relations with various concentrations and temperatures, with a view to investigate the exact nature of the molecular interaction. Adiabatic compressibility and intermolecular free length decreases with increase in concentration and temperature and gradually increases with concentration of solutes. These parameters have further used to interpret the hydrophilic part of the solute and molecular interactions in the solutions.

Keywords: Acoustic, thermodynamic parameters, electrolytes, molecular interaction, hydrophilic components of the mixtures.

1. Introduction

The ultrasonic velocity (u) , density (ρ) and viscosity (η)can be used to study the physicochemical behavior and molecular interactions such as ion-solvent interaction and solvent-solvent in pure liquids, liquid mixtures and solutions[1-3]. The nature and degree of molecular interactions in different solutions changes depending upon the nature of solvent, the structure of solute molecule and extent of solution taking place in the solution[4-5]. Acoustical impedance (z), adiabatic compressibility (β_a), intermolecular free length(L_f), relative association (R_A) are function of ultrasonic velocity .As these parameters throw more light on ion- ion and ion-solvent interactions, an ultrasonic and thermodynamic study of electrolytes.

NaCl and MgCl₂ having specialized properties like low melting but high boiling point. Both electrolytes are soluble in H₂O at 20^oC. Both electrolytes are insoluble in ether[6]. These electrolytes are also called as electrovalent compounds. NaCl is acts as a monovalent compound where MgCl₂ acts as a divalent compounds. As it occurs in metabolism of almost all living things, its interactions in an aqueous solution is of great value to the biological scientists.. It has wide applications in medicine and industry.

2. Experimental Section

All the chemicals used were of AR grade and dried over anhydrous CaCl₂ in desiccators before use. All solutions were prepared in deionized and distilled water (degassed by boiling), having specific conductivity ~ 10^{-6} S cm⁻¹. The stock solutions of 1M concentration were prepared by weighing the serine on a digital balance with an accuracy of $\pm 1 \times 10^{-4}$ g. Solutions of NaCl & MgCl₂ were made by mass on the mole fraction scale. Uncertainties in solution concentrations were estimated at $\pm 1 \times 10^{-5}$ mol kg⁻¹ in calculations. The solutions were kept in the special air tight bottles and were used within 12 hrs after preparation to minimize decomposition due to bacterial contamination. Ultrasonic velocity was measured with a single crystal interferometer (F- 81, Mittal Enterprises, New Delhi) at 2MHz The interferometer was calibrated against the ultrasonic velocity of water used at T = 303.15K. The present experimental value is 1508.80 ms⁻¹ which is in good agreement with literature value 1509.55 ms⁻¹. Accuracy in the velocity measurement was \pm 1.0 ms⁻¹. The density measurements were performed with recalibrated specific gravity bottle with an accuracy of $\pm 2x10^{-2}$ kg m⁻³. An average of triple measurements was taken into account. Sufficient care was taken to avoid any air bubble entrapment. Viscosity was measured with recalibrated Ostwald type viscometer. The flow of time was measured with a digital stop watch capable of registering time accurate to ± 0.1 s. An average of three or four sets of flow of times for each solution was taken for the purpose of calculation of viscosity. The accuracy of the viscosity measurements was \pm %. Accuracy in experimental temperature was 0.5 maintained at ± 0.1 K by means of thermostatic water bath.

3. Results and Discussion

From the measured values ultrasonic velocity(u), $density(\rho)$ and viscosity (η) various acoustical parameters such as the compressibility(β_a),acoustic impedance adiabatic (z). intermolecular free length (L_f) and relative association (R_A) were calculated by using the following relations [7,8,9,10]

Ultrasonic velocity (u) =
$$n \times \lambda$$
 (1)

Adiabatic compressibility (β_a) = $1/u^2 \rho$ (2)(3)

Acoustic impedance $(z) = u.\rho$ Intermolecular free length (L_f) = K / u.p^{1/2}

(4) Relative association (R_{A}) = (ρ / ρ_0). (u_o / u)^{1/3}

Where, K is the temperature dependant Jacobson constant¹¹. (5)

The values of K x 10^4 are taken as 207.556 x 10^{-8} ,209.431 x 10⁻⁸ and 211.306 x 10⁻⁸ at 303.15, 308.15 and 313.15K. T is the absolute temperature, ρ_0 , ρ and u_0 , u are the density and ultrasonic velocity of solvent and solution respectively.

International Journal of Science and Research (IJSR) ISSN (Online): 2319-7064, Impact Factor (2013): 4.438

www.ijsr.net

The experimentally measured values of ultrasonic velocity (u), density (ρ) and viscosity (η) of the solutions and calculated values of acoustical parameters such as adiabatic compressibility (β_a), acoustic impedance (z), intermolecular

free length (L_f) and relative association (R_A) are reported in Table -1 for the systems (water + NaCl) and Table-2 for the system (water+ $MgCl_2$) respectively.

Table 1: Ultrasonic velocities, densities, adiabatic compressibility's, acoustic impedances, free lengths ,relative association numbers of NaCl in double distilled water at 2MHz at 303.15K.

Double distilled Water +NaCl

m mol kg ⁻¹	u ms ⁻¹	ρ Kg m ⁻³	η Nm ⁻² s	βx10 ⁻¹⁰ m ² N ⁻¹	zx10 ⁻⁶ Nm ⁻²	${f L_f}{f A^0}$	R _A			
NaCl + Serine										
0.000	1551.80	1058.11	1.06195	3.92502	1.64180	0.41120	1.05285			
0.008	1564.88	1064.54	1.16268	3.83071	1.66816	0.40623	1.057 85			
0.017	1593.18	1078.71	1.37615	3.65472	1.71745	0.39679	1.06339			
0.026	1601.16	1081.52	1.55944	3.60831	1.73085	0.39426	1.06458			
0.034	1618.12	1087.01	1.61528	3.51368	1.75890	0.38905	1.06674			
0.043	1651.64	1094.01	1.87772	3.35085	1.80689	0.37994	1.06630			

 Table 2: Ultrasonic velocities, densities, adiabatic compressibility's, acoustic impedances, free lengths ,relative association numbers of MgCl₂ in double distilled water at 2MHz at 303.15K.

 Double distilled Water + MaCl

m mol kg ⁻¹	u ms ⁻¹	ρ Kg m ⁻³	η Nm ⁻² s	βx10 ⁻¹⁰ m ² N ⁻¹	zx10 ⁻⁶ Nm ⁻²	$\begin{array}{c} \mathbf{L_{f}} \\ \mathbf{A^{0}} \end{array}$	R _A		
$MgCl_2 + Serine$									
0.000	1560.51	1058.38	1.06103	3.88131	1.65101	0.40891	1.05089		
0.008	1581.00	1064.78	1.17284	3.74601	1.68851	0.40172	1.05622		
0.017	1599.76	1081.34	1.38746	3.61468	1.72934	0.39817	1.06489		
0.026	1618.88	1096.66	1.56448	3.48142	1.74234	0.39461	1.07540		
0.034	1627.20	1100.24	1.38931	3.43346	1.78992	0.38459	1.07749		
0.043	1642.80	1116.85	1.89523	3.32024	1.83336	0.38384	1.08969		

From the tables 1- and table -2, we observe the trends of all acoustical parameters with variation in concentration and temperature. The ultrasonic velocity increases with molal concentration of solute as well as rise in temperature. This increase in ultrasonic velocity in the aqueous solution of electrolytes may be attributed to the cohesion brought by the ionic hydration .The increase in density with molal concentration suggest a solute-solvent interaction exist between the electrolytes and water .In other words the increase in density may be interpreted to the structure maker of the solvent due to H-bonding[12,13]. The viscosity is an important parameter in understanding the structure as well as molecular interaction occurring in the solutions. From above tables, values of viscosities increase with concentration but decreases with temperature. These variations attributed to [14].The structural changes values of adiabatic compressibility (a) show decreasing trend with concentration as well as temperature which suggest the making and breaking of H-bonding. The intermolecular free length depends upon the intermolecular attractive and repulsive forces. Eyring and Kincaid[15] have proposed that L_f is a predominating factor in determining the variation of ultrasonic velocity of solution. The values of intermolecular free length listed in the table 1 & table2 which shows decreasing trend with concentration and temperature Hence it can be concluded that there is significant interaction between solute and solvent molecules due to which the structural arrangement is also affected. Thus it is clear from the above parameters that there is a strong association between water and electrolytes molecules showing hydrophilic nature[16].

4. Conclusion

A strong intermolecular H-bonding interaction exist between electrolytes and water. From the acoustical parameters it is concluded that H-bonding interaction is very strong at higher temperature and concentration.

5. Acknowledgement

One of the authors is thankful to shri .S.R. Kanhekar for helpful discussion.

References

- [1] Pal Amlendu ; Sureshkumar; *Ind J of Chem*, 44A, Dec**2005**, 2490-94.
- [2] SA Ikhe; ML Narwade; *Ind. J of Chem*, 44A, June **2005**, 1203-05.
- [3] Pankaj Atri; Madhusudan Reddy; P Venkatesu; *Ind J of Chem* May-June**2010** 736-42.
- [4] V kannappan; R. Palani; *.Ind J Chem*, secA,Jan**2007**, 54-59.
- [5] S Thirumaran; S Sudha; *J of Chem Pharm res*, **2010**(1) 327.
- [6] Sanjeevan Kharat; J.International Journal of Appl.Chem, Sept, 2008.Saravazyan A P ,Ann Rev Biophys Chem, 1991,20,1321.

International Symposium on Ultrasonics-2015, 22-24 January 2015

Department of Physics, Rashtrasant Tukdoji Maharaj Nagpur University, Nagpur, Maharashtra, India Licensed Under Creative Commons Attribution CC BY

- [7] S.P Naidu , P.Subrahmanyam & P Ravindra, J.of Pure and Appl. Ultrasn, 2001, 24, 1, 18–23.
- [8] A. Bahadur, M.S Shubha and K.C rao, *J. of Pure and Appl. Ultrason*, **2001**, 23, 28 -30.
- [9] D Das and D.K Hazra, *Indian J. of Physical Chemistry*, 2003, 77B, 5, 519 523.
- [10] T.Ramanjipa ., M.E Rao . & E Rajgopal, .*Indian J. of Pure & Appl. Physics*, **1993**, 31, 348 349.
- [11] Tamura , K Gekko , Biochemistry, 1995, 34, 1878-884,
- [12] VA Tabhane; OP Chimankar; S Manja; TK Naminarayanan; *Pure Appl Ultraso*, (21) 1999,67.
- [13] S Thirumaran; AN Kannapan; *Global J of Mole.sci*, 4(2) 2009 160.
- [14] PB Agrawal; Mohd Indrees Mohd Siddique; ML Narwade; *Ind J Chem*, 42A (5) 2003,1050.
- [15] B Eyring; JF Kincadid; ,J. Chem. Phys, 1938, 6,620.
- [16] VJ Rajendra; Pure Appl Ultrason.17 1995, 65.