

Comparative Effect of High-Intensity Interval Training and Continuous Aerobic Training on Cardiovascular Fitness and Fatigue Resistance in Competitive Football Players

Dr. Sarika Kukalore¹, Dr. Neeraj Singh²

¹Associate Professor, MPT Cardiothoracic, Department of Physiotherapy, Career College, Bhopal-462023, India

²Assistant Professor, MPT Sports, Department of Physiotherapy, Career College Bhopal-462023, India

Abstract: ***Background:** Structured cardiovascular endurance training has been shown to significantly improve aerobic capacity and recovery parameters in competitive football players. However, limited evidence exists comparing the relative effectiveness of high-intensity interval training (HIIT) and continuous aerobic training (CAT) on cardiovascular fitness and fatigue resistance in football-specific contexts. **Objective:** To compare the effects of HIIT and CAT on VO_{2max} , heart rate recovery, and fatigue resistance in competitive football players. **Methods:** Thirty competitive male football players (18–25 years) were randomly allocated into two groups: HIIT group ($n = 15$) and CAT group ($n = 15$). Both groups trained for 6 weeks, 4 sessions per week. VO_{2max} , heart rate recovery, and repeated sprint ability were assessed pre- and post-intervention. **Results:** Both groups demonstrated significant improvements in VO_{2max} and heart rate recovery ($p < 0.05$). However, the HIIT group showed significantly greater improvements in VO_{2max} and fatigue resistance compared to the CAT group ($p < 0.01$). **Conclusion:** While both training modalities enhance cardiovascular fitness, HIIT produces superior improvements in aerobic capacity and fatigue resistance in competitive football players.*

Keywords: HIIT, aerobic training, football, VO_{2max} , fatigue resistance

1. Introduction

Football performance depends on the ability to repeatedly perform high-intensity actions interspersed with brief recovery periods. Aerobic fitness plays a crucial role in sustaining performance, enhancing recovery, and delaying fatigue during match play.

Recent evidence, including structured endurance training interventions, has demonstrated significant improvements in aerobic capacity and cardiovascular efficiency in football players. However, endurance training methods vary considerably, with **HIIT** and **continuous aerobic training (CAT)** being the most commonly employed approaches.

HIIT is characterized by short bursts of high-intensity exercise followed by recovery periods, closely mimicking the physiological demands of football. In contrast, CAT involves prolonged exercise at moderate intensity and is traditionally used to enhance aerobic base fitness.

Given the differing physiological demands and adaptations associated with these methods, a direct comparison is essential to guide evidence-based conditioning practices in football.

2. Objectives

Primary Objective

- To compare the effects of HIIT and CAT on aerobic capacity (VO_{2max}) in competitive football players.

Secondary Objectives

- To compare changes in heart rate recovery between groups
- To evaluate fatigue resistance using repeated sprint performance

3. Research Methodology

Study Design

Randomized comparative experimental study.

Participants

Thirty competitive male football players aged 18–25 years.

Inclusion Criteria

- Minimum 2 years of competitive football experience
- Regular training ≥ 4 sessions/week
- Medically fit

Exclusion Criteria

- Recent musculoskeletal injury
- Participation in additional endurance programs

Intervention Protocol

Duration: 6 weeks

Frequency: 4 sessions/week

Group A – HIIT

- Intensity: 85–95% HRmax
- Protocol: 4–6 $\times$ 3-minute intervals
- Recovery: 2 minutes active recovery

Group B – Continuous Aerobic Training

- Intensity: 65–75% HRmax
- Duration: 30–40 minutes continuous running

Outcome Measures

- VO₂max
- Heart rate recovery (1 minute post-exercise)
- Repeated sprint ability (fatigue index)

Statistical Analysis


Paired t-test for within-group analysis and independent t-test for between-group comparison.



Significance set at **p < 0.05**.

4. Results

Table 1: Comparison of Cardiovascular Fitness and Fatigue Resistance parameters between HIIT and Continuous Aerobic Training Groups

Outcome Measure	HIIT Pre (mean± SD)	HIIT Post (mean± SD)	CAT Pre (mean± SD)	CAT Post (mean± SD)	p-Value (between groups)
VO ₂ max (ml.kg ⁻¹ .min ⁻¹)	48.5± 3.4	54.1± 3.6	48.3± 3.6	51.0± 3.5	<0.01**
Heart Rate Recovery (beats/ min)	25.1 ± 4.0	33.5± 4.3	24.8 ± 4.1	29.2 ± 4.0	<0.05*
Fatigue Index (%)	14.8 ± 2.1	10.2± 1.9	14.6 ± 2.3	12.9± 2.0	<0.01**





Both groups demonstrated significant improvements in $VO_2\max$ and heart rate recovery following training.

$VO_2\max$


- HIIT group showed significantly greater improvement compared to CAT ($p < 0.01$)

Heart Rate Recovery

- Both groups improved significantly
- Greater improvement observed in HIIT group ($p < 0.05$)

Fatigue Resistance

- HIIT group demonstrated significantly lower fatigue index values post-training ($p < 0.01$)



5. Discussion

The findings of this study indicate that both HIIT and continuous aerobic training are effective in improving cardiovascular fitness in competitive football players. However, HIIT resulted in superior improvements in VO_{2max} and fatigue resistance.

These results complement previous findings demonstrating the effectiveness of structured cardiovascular endurance training by further identifying **HIIT as a more efficient modality** for football-specific conditioning. The intermittent nature of HIIT closely replicates match demands, leading to enhanced physiological adaptations relevant to football performance.

6. Conclusion

HIIT is more effective than continuous aerobic training in improving aerobic capacity, heart rate recovery, and fatigue resistance in competitive football players. Conditioning programs should prioritize high-intensity interval-based strategies for optimal performance enhancement.

7. Clinical Implications

- Guides sports physiotherapists and conditioning coaches
- Supports HIIT-based football conditioning
- Enhances match-specific endurance and recovery

8. Limitations

- Small sample size
- Short intervention duration
- Male participants only

9. Future Recommendations

- Inclusion of female athletes
- Longitudinal studies
- Integration with neuromuscular and strength training

References

- [1] Bangsbo J, Mohr M, Krstrup P. Physical and metabolic demands of training and match-play in the elite football player. *J Sports Sci.* 2006;24(7):665–674.
- [2] Helgerud J, Engen LC, Wisloff U, Hoff J. Aerobic endurance training improves soccer performance. *Med Sci Sports Exerc.* 2001;33(11):1925–1931.
- [3] Buchheit M, Laursen PB. High-intensity interval training, solutions to the programming puzzle. *Sports Med.* 2013;43(5):313–338.
- [4] Iaia FM, Bangsbo J. Speed endurance training is a powerful stimulus for physiological adaptations and performance improvements of athletes. *Scand J Med Sci Sports.* 2010;20(Suppl 2):11–23.
- [5] Impellizzeri FM, Rampinini E, Castagna C, Bishop D, Ferrari Bravo D, Tibaudi A, et al. Validity of a repeated-sprint test for football. *Int J Sports Med.* 2008;29(11):899–905.
- [6] Hoff J, Wisloff U, Engen LC, Kemi OJ, Helgerud J. Soccer specific aerobic endurance training. *Br J Sports Med.* 2002;36(3):218–221.
- [7] Buchheit M, Gindre C. Cardiac parasympathetic regulation: respective associations with cardiorespiratory fitness and training load. *Am J Physiol Heart Circ Physiol.* 2006;291(1):H451–H458.
- [8] Midgley AW, McNaughton LR, Jones AM. Training to enhance the physiological determinants of long-distance running performance. *Sports Med.* 2007;37(10):857–880.
- [9] Weston M, Taylor KL, Batterham AM, Hopkins WG. Effects of low-volume high-intensity interval training

- (HIT) on fitness in adults: a meta-analysis of controlled and non-controlled trials. *Sports Med.* 2014;44(7):1005–1017.
- [10] Dupont G, Akakpo K, Berthoin S. The effect of high-intensity interval training on performance in elite soccer players. *J Strength Cond Res.* 2004;18(3):584–589.
- [11] Krstrup P, Mohr M, Ellingsgaard H, Bangsbo J. Physical demands during an elite female soccer game: importance of training status. *Med Sci Sports Exerc.* 2005;37(7):1242–1248.
- [12] Laursen PB, Jenkins DG. The scientific basis for high-intensity interval training: optimising training programmes and maximising performance in highly trained endurance athletes. *Sports Med.* 2002;32(1):53–73.